


ASTRONOMICAL LESSONS - GORE.

# ASTRONOMICAL LESSONS



J. E. GORE







758

J. B. Dabary  
11 Sep 1893.

ASTRONOMICAL LESSONS.

OTHER WORKS BY THE SAME AUTHOR.

*In Crown 8vo, cloth extra. Price 10s. 6d. post free.*

**THE SCENERY OF THE  
HEAVENS:**

A POPULAR ACCOUNT OF ASTRONOMICAL  
WONDERS.

\* \* This beautiful work is illustrated by photographs of  
Star Clusters and Nebulae from original photographs  
taken at the Paris Observatory, and by Mr. ISAAC  
ROBERTS at Liverpool; also drawings from recent  
sketches by well-known Astronomers.

*In Crown 8vo, beautifully illustrated and appropriately  
bound in cloth extra. Price 7s. 6d. post free.*

**PLANETARY AND STELLAR  
STUDIES;**

OR, SHORT PAPERS ON THE PLANETS,  
STARS, AND NEBULÆ.

\* \* Detailed List with Press Opinions post free.

LONDON: SUTTON, DROWLEY & CO.

(Amalgamation of R. SUTTON & CO. and ROPER & DROWLEY),

11, LUDGATE HILL, E.C.






FIG. 1.—MAN-OF-WAR "HULL DOWN" AS SEEN WITH A TELESCOPE. [Frontispiece.]

# ASTRONOMICAL LESSONS ;

OR,

## CHAPTERS ON THE ELEMENTARY PRINCIPLES AND FACTS OF ASTRONOMY

FOR THE USE OF STUDENTS AND YOUNG PEOPLE.

BY

JOHN ELLARD GORE,

F.R.A.S., M.R.I.A., F.R.G.S.I., ASSOC.M.I.C.E.,

*Honorary Member of the Liverpool Astronomical Society, the "Arts Society,"  
and "The British and Foreign Association."*

AUTHOR OF

"The Scenery of the Heavens," "Planetary and Stellar Studies,"  
"Southern Stellar Objects," &c.

London :

SUTTON, DROWLEY, & CO.,

11, LUDGATE HILL, E.C.

1890.

[All rights reserved.]





FIG. 1.—MAN-OF-WAR "HULL DOWN" AS SEEN WITH A TELESCOPE. [*Frontispiece.*]

# ASTRONOMICAL LESSONS ;

OR,

CHAPTERS ON THE ELEMENTARY PRINCIPLES  
AND FACTS OF ASTRONOMY

FOR THE USE OF STUDENTS AND YOUNG PEOPLE.

BY

JOHN ELLARD GORE,

F.R.A.S., M.R.I.A., F.R.G.S.I., ASSOC.M.I.C.E.,

*Honorary Member of the Liverpool Astronomical Society, the "Arts Society,"  
and "The British and Foreign Association."*

AUTHOR OF

"The Scenery of the Heavens," "Planetary and Stellar Studies,"  
"Southern Stellar Objects," &c.

London:

SUTTON, DROWLEY, & CO.,

11, LUDGATE HILL, E.C.

1890.

[*All rights reserved.*]



"The works of the Lord are great, sought out of all them that have  
pleasure therein."—Psa. cxi. 2.

"Snatch me to heaven ; thy rolling wonders there,  
World beyond world, in infinite extent,  
Profusely scatter'd o'er the blue expanse  
Show me ; their motions, periods, and their laws  
Give me to scan."

THOMSON (*Autumn*).



## PREFACE.

---

THE following chapters on the elementary principles and facts of Astronomy are intended as an introduction to larger and more advanced works on the subject, and will, it is hoped, be found useful to young people and beginners in the study of this fascinating science

*October, 1890.*

J. E. G.



## CONTENTS.

---

CHAP.	PAGE
I. THE FIGURE OF THE EARTH ... ..	15
II. THE DIURNAL MOTION OF THE EARTH ...	20
III. THE FIXED STARS AND THE SUN'S APPARENT MOTION ... ..	25
IV. THE ANNUAL MOTION OF THE EARTH AND THE SEASONS ... ..	40
V. THE EQUATION OF TIME AND LEAP YEAR ...	53
VI. THE SUN ... ..	57
VII. THE MOON ... ..	61
VIII. ECLIPSES ... ..	72
IX. MERCURY ... ..	79
X. VENUS ... ..	83
XI. MARS ... ..	93
XII. THE MINOR PLANETS ... ..	96

CHAP.	PAGE
XIII. JUPITER ... ..	98
XIV. SATURN ... ..	104
XV. URANUS AND NEPTUNE ... ..	108
XVI. COMETS ... ..	113
XVII. DOUBLE AND BINARY STARS ... ..	116
XVIII. VARIABLE STARS ... ..	125
XIX. THE NEBULÆ ... ..	130

## LIST OF ILLUSTRATIONS.

FIG.	PAGE
1.—MAN-OF-WAR "HULL DOWN" AS SEEN WITH A TELESCOPE ... ..	<i>Frontispiece</i>
2.—IMAGE OF THE SETTING SUN REFLECTED FROM A CALM SEA ... ..	17
3.—SENSIBLE AND RATIONAL HORIZON ... ..	22
4.—THE PLOUGH AND THE POLE STAR ... ..	31
5.— ... ..	35
6.— ... ..	37
7.—MOTION IN AN ELLIPTIC ORBIT ... ..	45
8.—THE SEASONS ... ..	49
9.—MOTION OF THE MOON ... ..	62
10.—PHASES OF THE MOON ... ..	64
11.—ROTATION OF THE MOON ON ITS AXIS ... ..	67
12.) 13.) —TO ILLUSTRATE PHENOMENON OF HARVEST MOON ... ..	69



FIG.	PAGE
14.—ECLIPSE OF THE MOON ... ..	75
15.—RELATIVE POSITIONS OF THE EARTH AND VENUS FROM GREATEST "ELONGATION" OF VENUS, THROUGH INFERIOR CONJUNCTION ...	87
16.—TRANSIT OF VENUS ... ..	89
17.—RELATIVE POSITIONS OF THE EARTH AND MARS BEFORE AND AFTER "OPPOSITION" OF MARS	93
18.—ECLIPSES OF JUPITER'S SATELLITES, AS SEEN FROM THE EARTH ... ..	101
19.—POSITION OF NEPTUNE OBSERVED BY LALANDE MAY 8TH AND 10TH, 1795 ... ..	111
20.—APPARENT ORBIT OF THE COMPANION TO SIRIUS AS COMPUTED BY J. E. GORE ...	121
21.—THE VICINITY OF MIRA ( $\alpha$ ) CETI ... ..	127
22.—THE VICINITY OF ALGOL ( $\beta$ PERSEI) ...	129

## CHAPTER I.

## THE FIGURE OF THE EARTH.

THE word Astronomy means the science which treats of the heavenly bodies, their motions, and the laws which govern them. It is derived from two Greek words, *αστηρ* (*aster*) a star, and *νομος* (*nomos*) a law—the law of the stars. It is a most interesting study, and has attracted the attention of mankind since the earliest ages of antiquity. We will commence our study with the figure and dimensions of the earth on which we live. The shape of the earth is nearly that of a globe or sphere.

The reader may ask: How can it be proved that the earth is a globe, and not a plane surface, as ordinary observation would lead us to suppose?

There are several convincing proofs of the globular shape of the earth. For instance, if the earth were a flat plane, when the sun rose above the horizon of any place it would at once illuminate the whole terrestrial surface. This, we know, is not the fact, for the sun rises in London several hours before it rises in New York. How the supporters of the

"flat earth" theory explain this admitted fact I do not know. It certainly could not be explained on any common-sense principle. But perhaps the most familiar, and certainly the most convincing proof is the appearance of ships at sea. If you stand on the shore on a calm day, and watch a vessel sailing away from your eye, would you not expect to see the masts disappear before the hull, if the surface of the sea were a level plane? The hull, being so much larger than a comparatively thin mast, should remain for a longer time visible.

You will see, however, that the very reverse of this takes place. The hull of the ship is the *first* portion to disappear, and then the masts gradually sink out of sight, till at last nothing but the extreme top of the highest mast remains visible. This clearly shows that the surface of the sea is globular, and hides the ship's hull from our view. You will see also that the bounding-line of the horizon, called the *offing* by sailors, is a sharp, clearly defined line, not fading away in haze and distance, as would be the case were the surface of the water a level plane.

This is the appearance as seen with the naked eye, but—it may be asked—if we examined the ship with a telescope, would not the hull then become visible? No; the appearance would be exactly the same. The frontispiece represents the telescopic view of a "man of war"<sup>1</sup> when it is "hull down," as the sailors term it. Were the earth flat, as some

<sup>1</sup> The *Warspite*.

mistaken people imagine, it is quite impossible that any such appearance could ever be seen. The ship would continue visible as a *whole*, and would become apparently smaller and smaller, until at last it became invisible even in a powerful telescope. This seems quite evident, but there are other proofs of the earth's rotundity.

The shadow of the earth on the moon during a lunar eclipse is another proof. An eclipse of the



FIG. 2.—IMAGE OF THE SETTING SUN REFLECTED FROM A CALM SEA.

moon is caused by the earth passing between the sun and the moon, and in a partial eclipse we see that the shadow on the moon's face is bounded by a circular line. Now nothing but a spherical body could, in *all* positions, throw a circular shadow. A very interesting proof of the globular form of the earth has recently been given by Prof. Ricco, the well-known astronomer of the Palermo Observatory. He finds that the image of the sun, reflected from the surface



of a calm sea, is of a well-marked oval shape, and he expresses his surprise that the ancient astronomers did not notice this phenomenon, and infer from it that the earth is really globular. M. Ricco made his observations with a telescope of  $4\frac{1}{2}$  inches aperture, from a height of 236 feet above the sea level, and about  $1\frac{1}{4}$  miles from the shore. He finds that when the sun's lower limb is touching the sea horizon, the vertical diameter of the reflected image is reduced about ten minutes of arc, or nearly one-third of the sun's apparent diameter (see fig. 2). This result agrees with theory, and *proves* that the earth is a sphere.

The earth is, however, not a perfect sphere. It is slightly flattened at the north and south poles—somewhat similar to an orange (but not to so great an extent); the diameter at the equator being about  $26\frac{1}{2}$  miles longer than that through the poles.

If you look at an artificial globe you will see that it is mounted on an axis, on which it turns. This represents the earth's axis, and the extremities of this axis are termed the poles. The earth of course has no real axis, like the artificial globe, but as it turns round once in the 24 hours, astronomers call the imaginary line round which it rotates the *axis*, and the extremities of this line are the earth's *poles*. This imaginary axis points very nearly to a tolerably bright star, known to astronomers as the Pole Star.

The equator is an imaginary circle round the earth,

every point on which is at an equal distance from the poles. The most accurate measurements make the diameter at the equator  $7925\frac{1}{2}$  miles, and at the poles 7899 miles, the difference being, as I have said, about  $26\frac{1}{2}$  miles. The mean or average diameter is therefore  $7912\frac{1}{4}$  miles.

## CHAPTER II.

## THE DIURNAL MOTION OF THE EARTH.

I SPOKE of the rotation of the earth on its axis in 24 hours. This is the cause of day and night. But—it may be asked—if the earth really rotated on its axis should not the motion be, in some way, visible to observation? So it is; in the apparent motion of the sun, moon, and stars. In what other way the motion *could* become sensible I cannot conceive. It must be remembered that the motion is very steady and uniform, that the earth's atmosphere is carried round with it, and that we have no external objects, except the heavenly bodies, to which we can refer the motion. When on board ship, out of sight of land, with a cloudy sky, and neither sun, moon, or stars visible, it is impossible to tell (without watching the "wake" of the ship in the water) whether she is moving in a straight line, or going round in a circle. But if you were near land, and looking out of the cabin window, you could at once detect the motion of the vessel by watching objects on the shore, which would seem to move in the opposite direction. In

the same way when travelling in a fast railway train, you see outside objects apparently flying the other way. We must therefore conclude that either the earth turns on its axis, or the whole star sphere rotates round the earth every 24 hours; which is the more probable? there can be no third alternative. It certainly seems more probable that a comparatively small body, like the earth, should turn round on its axis than that the gigantic sphere of stars should rotate. The reader may ask, however: Is there any *direct proof* of the earth's rotation?

Yes; an experiment devised by M. Foucault proves conclusively that the earth *does* rotate on its axis in 24 hours. If a heavy weight be suspended by a long wire, and caused to vibrate like a pendulum in a certain direction, say north and south, it will be found, if attentively watched, that the plane of vibration rotates. At the equator there would be no effect, but at the north and south poles the plane of vibration would make one rotation in 24 hours; or in 6 hours from the commencement of the experiment, the pendulum would be vibrating from east to west, instead of from north to south. We cannot, of course make the experiments at the poles, but experiments made at Paris agree with the mathematical theory of the motion, and prove, beyond all doubt, that the earth *does* rotate on its axis.

Here is another proof: As the top of a high tower is further from the earth's centre than the base of the tower, if the earth rotates the top will move faster



than the base. Hence if a ball be dropped from the top, we should expect it to fall a little to the east of the vertical line. Careful experiments have shown that such is the case. There are some other proofs,<sup>\*</sup> but the two I have just given are quite sufficient to establish the truth of the theory to any reasoning mind.

In speaking of the horizon it should be mentioned that there is a *sensible* and a *rational* horizon. The




FIG. 3.—SENSIBLE AND RATIONAL HORIZON.

*sensible* horizon is the circle bounding the view of a spectator from any point on the earth's surface. The *rational* horizon is that supposed to be seen from the centre of the earth. In the accompanying figure APB is the *sensible* horizon of the point P, and DCE is the *rational* horizon of the same point. From a high point we should, however, see a greater

<sup>\*</sup> We know that Mars, Jupiter, and Saturn rotate on their axes (and probably all the other planets). It seems, therefore, from analogy, highly improbable that the earth should be the only planet having no axial rotation.

distance. From the elevation Q (see fig. 3) the distance of the *sensible* horizon would be QS or QT, and the higher the point of observation, the greater the distance visible. The *rational* horizon is, of course, an imaginary one, so that all the motions of the heavenly bodies must be referred to the *sensible* horizon.

The setting sun remains visible for a longer time as seen from the top of a mountain than from the plain below. Some years since an aeronaut ascended in a balloon from Dublin, and was carried across the Irish Channel. When approaching the coast of Wales the balloon descended, a little after sunset, nearly to the surface of the sea. The aeronaut then threw out nearly all his ballast, and the balloon suddenly rising to a great height the sun again became visible, producing the curious phenomenon of a sunrise in the west!

The difference between the *sensible* and the *rational* horizons will not, however, make any difference in the extent of the heavens visible to a spectator on the surface of the earth. For, as the earth is a mere point in comparison with the nearest of the stars, there will be no perceptible difference whatever between the heavens as seen from the surface and the imaginary view visible from the centre of the earth.

Thus we always see the *whole* hemisphere; and about half the number of stars in the heavens are visible at one time. If you imagine the small point representing the centre of the circle in fig. 3 to represent the

earth you will see that a whole hemisphere is visible from every point on the earth's surface. From a high mountain, or from a balloon, slightly *more* than a hemisphere would be visible. It is true that when clouded the sky looks somewhat like a flattened arch, but this is clearly due to the fact that the clouds overhead are really nearer to the eye than those near the horizon (the height of the clouds at both points being the same). In a cloudless sky, however, the hemispherical shape is very evident and unmistakable.

### CHAPTER III.

#### THE FIXED STARS AND THE SUN'S APPARENT MOTION.

CATALOGUES of the stars have been made by astronomers, even of thousands not visible to the naked eye. But, the reader may ask, is this possible? the number visible even to the naked eye seems quite innumerable.

This is, however, a popular fallacy—due chiefly to an optical illusion. But in reality the number of stars distinctly visible to the naked eye is comparatively *small*. The number given in the catalogues of the ancient astronomers is *very* small. That of Hipparchus contains only 1,022, and that of Hevelius—the last of the astronomers who observed the places of stars without a telescope — only 1,533, observed at Dantzic.

The number really visible to good eyesight is, however, somewhat larger. Those visible without optical aid have been carefully mapped by two eminent astronomers, Behrmann and Heis, both of whom possessed exceptionally keen eyesight. For the stars visible in northern latitudes Heis found a total of



5,356, and Behrmann for the southern portion (not observed by Heis) 2,306, making a total for the whole heavens in both hemispheres of 7,662. Of course only one-half this number (one hemisphere) can be seen at any one time, or an average of about 3,830.

As many of the stars observed by Heis and Behrmann are very faint, even this small number would not be visible to every one. If we confine the enumeration to stars of the 6th magnitude—about the average limit of most eyes—the total, in both hemispheres, is about 4,000, giving an average of only 2,000 as visible at any one time to ordinary eyesight.

From these figures it will be seen that the lucid stars (as those visible to the naked eye are called) are very thinly scattered over the surface of the heavens. Were there on an average even one star to each space of the area of the full moon there would be no less than 180,000 stars visible to the unassisted eyesight, or 90,000 at one time. This will give an idea of how very sparsely scattered the lucid stars really are. This fact may appear very surprising. Some may think that millions are visible on a clear night. The idea, indeed, seems a very common one, but it is none the less erroneous. The student may depend upon it that astronomers do not make serious mistakes of this kind. The numerical results I have given are such that it would be in vain to attempt to disprove them. Any one who tries to do so merely shows his

ignorance and inability to understand scientific facts and reasoning.

The number of stars is, however, largely increased when the heavens are examined with a telescope. Even with a good opera-glass the number is considerably increased, and the more powerful the telescope the larger the number visible. Not, however, in proportion to the size of the telescope. The number of telescopic stars appears to be limited also. Assuming that an infinite number of stars were scattered through infinite space, it may be *proved* mathematically that the whole heavens would shine with the brightness of the sun! (and with a corresponding amount of heat!) But even on a clear night the sky is more remarkable for its blackness than for its light. This is another surprising result, but it cannot be disputed.

To give an idea of the way in which the stars apparently "thin out" as we examine the fainter ones, I may mention that an estimate has been made of the total number of stars visible with the giant telescope of the Lick Observatory, California—recently constructed—and two of the greatest American astronomers, Professors Newcomb and Young, give as the probable number 100 millions. Now on the theory of uniform distribution through infinite space, I find the number which *should* be visible in this great telescope is over 3,000 millions, or 30 times the number which it actually shows!

How is this very limited number of stars to be explained? Some astronomers have suggested that

the luminiferous ether of space may—for stars at a vast distance—absorb the light of the fainter stars altogether. But if we are to consider the ether as a *perfect* fluid, as scientific men maintain, this explanation will not hold good. The most probable explanation seems to be that the sun and Solar System belong to a cluster of limited dimensions, in fact that the *visible* universe is really limited. If other external universes exist—as is probably the case, space being infinite—they are possibly invisible to us owing to immensity of distance.

The student will notice that the stars are of very different degrees of brightness. St. Paul says, "One star differeth from another star in glory." Those visible to the naked eye have been divided—even by the ancient astronomers—into six magnitudes, the brightest being called first magnitude, those clearly fainter second magnitude, and so on to the faintest visible to ordinary eyesight, which are called sixth magnitude. Telescopes, however, continue this series of magnitudes further, and with the largest telescopes the faintest stars are considered to be of the 16th and even 17th magnitude.

It will also be noticed that among the brightest stars some are perceptibly brighter than others. The earlier astronomers included all the brightest stars in the first magnitude, but Sirius, for instance, is far brighter than any of the others; modern observations show it to be nearly 2 magnitudes brighter than an average star of the first magnitude. The bright stars

Arcturus, Capella, Vega, and Procyon in the northern hemisphere, and Canopus,  $\alpha$  Centauri, and Rigel in the southern are also distinctly above the average.

The following stars may be considered as average stars of each magnitude:—Of the 1st magnitude, Aldebaran, Altair, and Spica; of the 2nd magnitude,  $\alpha$  Persei,  $\alpha$  Andromedæ, and the Pole Star in the northern hemisphere, and  $\beta$  Argûs,  $\lambda$  Scorpii, and  $\gamma$  Crucis in the southern hemisphere. Of the 3rd magnitude,  $\beta$  and  $\delta$  Draconis,  $\zeta$  Tauri,  $\epsilon$  Virginis,  $\beta$  Cygni,  $\theta$  Aquilæ, and  $\gamma$  Gruis. Of course between each magnitude there are a number of stars of intermediate brightness, the heavens containing stars of all grades of brilliancy, from Sirius to the faintest star visible in the Lick telescope.

The names just mentioned refer to the imaginary constellation figures into which the ancients grouped the stars for convenience of reference. These groups have been retained, with few alterations, by modern astronomers. The Greek letters were added by Bayer in 1603.

Before attempting to identify these constellations in the sky, we must know how to find the four points of the compass, North, South, East, and West. The student will know that at noon the Sun is due South, and that looking at it then, the North is behind him, the East on his left hand, and the West on his right hand.

But at night—it may be asked—when we have not the sun to guide us, how are we to find the north?



The student will probably have heard of the Pole Star. He may also know the seven bright stars forming the Plough, or Charles' Wain. Here is a drawing of it (fig. 4). It is in the constellation of *Ursa Major*, or the Great Bear. Well, a line drawn from  $\beta$  to  $\alpha$ , and produced, will nearly pass through the Pole Star. Not *exactly*, but near enough for the purpose of identifying the Pole Star, which, as I have said, is a standard star of the second magnitude, and therefore a fairly conspicuous one. For this reason these two stars are called the "pointers."

The Pole Star is not *exactly* at the Pole, but so near it that to *ordinary* observation it does not perceptibly alter its position. This star will be always useful in finding the north, and from it, and the stars of the Plough, the positions of other stars may easily be determined. For instance, if the curve formed by the three stars in the tail of the Bear ( $\epsilon$ ,  $\zeta$ ,  $\eta$  in the diagram) is continued on, it will pass near a very bright star. This is Arcturus, one of the brightest stars visible in Europe. Again, if we draw an imaginary line from  $\gamma$  to  $\beta$  and produce it, it will pass near another very bright star. This is Capella, or  $\alpha$  Aurigæ, the principal star of the constellation Auriga, the Charioteer. Again, if we draw a line from  $\delta$  to  $\beta$  and produce it, it will pass near two tolerably bright stars. These are Castor and Pollux in Gemini or the Twins, the northern of the two being Castor, and the lower, and somewhat the brighter of the two, Pollux. Another line from  $\delta$  to  $\gamma$  will pass




FIG. 4.—THE PLOUGH AND THE POLE STAR.

near a bright star called Regulus, or  $\alpha$  Leonis, sometimes called *Cor Leonis*, or the Lion's Heart. This star is in the handle of the so-called "Sickle," the figure formed by  $\alpha$ ,  $\eta$ ,  $\gamma$ ,  $\zeta$ ,  $\mu$ , and  $\epsilon$  in Leo, which very much resembles a reaping hook.

On the opposite side of the Pole Star from the Plough you will see five conspicuous stars, forming a W-shaped figure. This is Cassiopeia's Chair. In this way, and with the help of a celestial globe or star atlas, you will soon learn to know all the brightest stars at a glance, and from these brighter stars the fainter ones can be found when required. The star Regulus is remarkable for another reason : it lies close to the ecliptic.

The ecliptic is an imaginary circle in the sky which the sun apparently moves along during the course of the year. The term ecliptic is derived from the fact that eclipses of the sun and moon occur in or near this circle.

The sun, however, seems to have a motion across the heavens every day. This, called the *diurnal* motion, is simply due to the rotation of the earth on its axis once in 24 hours. This apparent motion, which is familiar to every one—even to those ignorant of astronomy—is quite distinct from the *annual* motion, which can only be detected by careful observation.

In what way can we detect this annual motion of the sun? If we take any well-known star near the ecliptic, and visible towards the west after sunset,

and if we observe this star every evening for some weeks, we shall soon notice that it sets earlier each evening, and at last approaches so near the sun that it is lost in his rays, and can no longer be seen. After some weeks this same star will become visible in the morning sky before sunrise, showing that it has apparently passed behind the sun. It then rises earlier and earlier each morning until it rises at sunset, when it is said to be in "opposition" (or opposite to) the sun, and is then visible all through the night for several months, until at last it again rises and sets with the sun.

This apparent motion of the sun among the fixed stars is due to the *real* motion of the earth round the sun, and when once recognized seems quite as obvious as the diurnal motion.

The student may ask, Do *all* the stars disappear and reappear in this way? Well, most of them do, at least in northern and southern latitudes. Those, however, within a certain distance of the Pole—equal to its altitude (or height above the horizon of the place)—never rise or set, as they do not pass below the horizon at any time. These are called *circumpolar* stars, and are visible at all seasons of the year, and at all times of the night.

The altitude of the Pole Star—or rather the altitude of the Pole—is its angular elevation above the horizon, and is equal to the latitude of the place of observation. The proof of this is very simple, and from the student's knowledge of Euclid he will



easily understand it. Let APEBD (fig. 5) represent the earth, C its centre, and AB its axis, pointing towards the pole in the direction AS. Let DCE, at right angles to AB, represent the earth's equator. Let P be the place of observation, and FPG the *sensible* horizon of the place—at right angles to PC. Then, since the Pole Star is at an immense distance from the earth, the line PT pointing to the Pole is parallel to CAS. Hence the angle TPF is equal to the angle PFC. But since the angles FPC and ACE are right angles, we have PFC and FCP together equal FCP and PCE. Take away the common angle FCP, and PFC is equal to PCE. But PCE is the latitude of the place P, being the angular distance from the equator E. Therefore PFC, or TPF, is the latitude of the place P. This relation is very useful for finding the latitude at sea by measurement of the sun's altitude at noon.

Before explaining the method of finding the latitude I must first explain that there is another imaginary circle in the heavens called the *celestial equator*. This is the circle in which the plane of the earth's equator, if produced, would intersect the star sphere. Hence, at the poles of the earth this circle will coincide with the horizon, and at the equator it will pass through the zenith, or point exactly overhead. The plane of the ecliptic is inclined to the plane of the equator at an angle of  $23\frac{1}{2}$  degrees, this being the angle between the earth's polar axis and a perpendicular to the plane of its orbit, or the plane




FIG. 5.

of the ecliptic.<sup>1</sup> See fig. 6, in which NESQ represents the earth, N the North Pole, S the South Pole, EQ the earth's equator, AB the plane of the ecliptic. It will be seen that the plane of the ecliptic is at one point raised  $23\frac{1}{2}$  degrees above the equator, and at the opposite point is  $23\frac{1}{2}$  degrees below it. Now the sun in its apparent motion round the heavens and moving along the ecliptic will at one point be situated  $23\frac{1}{2}^{\circ}$  north of the celestial equator. This occurs on the 21st of June, the "longest" day. Six months afterwards, or on the 21st of December, the "shortest" day, it will be  $23\frac{1}{2}^{\circ}$  south of the equator, and at two other points it will be exactly on the equator. These points, where the plane of the ecliptic intersects the plane of the equator, are called the *equinoctial* points, because day and night are then of equal length. The sun arrives at these points on March 21st, the *Vernal Equinox*, and on Sept. 21st, the *Autumnal Equinox*. At all other periods of the year it is either above or below the equator, and its distance north or south of this line—technically called its *Declination*—is given for each day of the year in a book called the *Nautical Almanack*, which is published some years in advance for the use of vessels making long voyages.

We know that every day at noon the sun is due south, and at its highest point above the horizon—or on the meridian of the place, as astronomers term it.

<sup>1</sup> The inclination of the polar axis to the plane of the ecliptic is of course  $90^{\circ} - 23\frac{1}{2}^{\circ} = 66\frac{1}{2}^{\circ}$ .




FIG. 6.



Now, to find his latitude at sea the captain of a ship measures the altitude of the sun at noon with an instrument called a sextant, and then knowing from the Nautical Almanack the sun's declination on that particular day, he can find the altitude of the celestial equator, and thence the latitude of the place of observation. There are some small corrections to be made to the observations, but I will explain the general principles of the method. If the sun is at the time north of the equator the captain deducts the declination from the observed altitude, and if south of the equator he adds it. He then subtracts the altitude from 90 degrees, and the difference is the latitude of the place. Let us try an example. The altitude of the sun at noon on the 21st of June is observed from a ship to be  $60^{\circ}$ ; what is the latitude? On the 21st of June the sun's declination is  $23\frac{1}{2}^{\circ}$  north of the equator. This must be deducted from  $60^{\circ}$ , which gives  $36\frac{1}{2}^{\circ}$  for the altitude of the equator, and  $36\frac{1}{2}^{\circ}$  deducted from  $90^{\circ}$  gives  $53\frac{1}{2}^{\circ}$ , the required latitude.

If the same altitude  $60^{\circ}$  was observed on the 21st of December, what would then be the latitude? In this case  $23\frac{1}{2}^{\circ}$  must be *added* to  $60^{\circ}$ , which gives  $83\frac{1}{2}^{\circ}$ , and this deducted from  $90^{\circ}$  leaves  $6\frac{1}{2}^{\circ}$  for the latitude.

Now, supposing the observed altitude on the shortest day was found to be  $70^{\circ}$  instead of  $60^{\circ}$ , let us find the latitude.  $70^{\circ}$  added to  $23\frac{1}{2}^{\circ}$  gives  $93\frac{1}{2}^{\circ}$ . But how can this be deducted from  $90^{\circ}$ , as it is a greater number? Well, the difference would be

in algebra *minus*  $3\frac{1}{2}$  degrees, or  $3\frac{1}{2}$  degrees less than zero. This shows that the celestial equator is  $3\frac{1}{2}$  degrees to the north of the zenith, from which we conclude that the place of observation is *south* of the terrestrial equator, or in  $3\frac{1}{2}$  degrees *south* latitude.

The latitude could also be found by observing at night the meridian altitude of a star whose position is known with reference to the celestial equator; but at sea it is usually found more convenient to observe the sun.

I have explained how the latitude of a place is found. How can the longitude be determined? This is a more difficult problem than the other. There are several methods of finding the longitude, but the one generally used at sea is the method by "lunar distances." In the Nautical Almanack the angular distance of the moon's centre from certain bright stars and planets is given for every three hours. If, then, the distance of the moon from a bright star or planet is measured with the sextant at any time, the Greenwich mean time of the observation can be calculated, and from the local time at the place the longitude may be found, reckoning 15 degrees of longitude to each hour of time. The local time may be found from sextant observations of the sun's altitude.

## CHAPTER IV.

THE ANNUAL MOTION OF THE EARTH AND  
THE SEASONS.

IN addition to the diurnal motion of the earth on its axis the earth has another motion—an annual motion round the sun. This motion causes the different seasons of the year.

But before I explain how the seasons are due to the earth's annual motion, perhaps the student would like to hear some proof that such a motion actually takes place. For it may seem to him that the sun's apparent motion among the stars might be explained equally well by a real motion of the sun round the earth. And he would be quite correct in that idea; the observed motion could be equally well explained on either supposition. But the evidence in favour of the earth's motion is so great that it seems impossible to resist it. As I explained, in speaking of the earth's rotation on its axis, we could not expect to be in any way conscious of its motion through space in its journey round the sun. The motion is so smooth and uniform that we could have no possible means of

ascertaining that such a motion exists, except by watching some *external* object, and such an apparent motion we see in the sun. We have now to determine whether it is the sun or the earth which performs a revolution in the course of a year. One or other of these motions *must* take place; there is no other alternative.<sup>1</sup> The best proofs of the earth's motion round the sun are mathematical ones, and for this reason, probably, the theory of the earth's motion is disputed by paradoxers, who are unable to understand the mathematical reasoning on which it is based. One proof is as follows :—The sun's distance from the earth can easily be ascertained by astronomical observations. This distance has been very approximately determined, and admits of no reasonable doubt. Knowing this distance and the sun's apparent diameter, its actual size follows by a very simple calculation. In this way it has been found that the sun is a vastly larger body than the earth. Now Sir Isaac Newton has proved that when two bodies revolve round each other, according to the law of gravitation, they must both revolve round a point between them, which is called "the common centre of gravity." In the case of the sun and earth the centre of gravity—owing to the vast size of the sun—is situated near the centre of the sun, and hence,

<sup>1</sup> The idea that the sun revolves round the earth once in 24 hours, as maintained by the supporters of the "flat earth" theory, is altogether preposterous and unworthy of serious consideration.



according to this principle, the earth must revolve round that point, the sun remaining nearly stationary.

Suppose a large globe and a small one are connected by a rod, and we try to balance the combination on another rod, it will be found that the point at which both globes will balance is very much nearer to the larger globe than to the smaller one, and the greater the difference in the size of the globes the nearer will this point fall to the larger globe. This point is called the "centre of gravity" of the two globes. If the larger globe is very large in comparison with the small one it will be found that the point falls within the surface of the larger globe, as happens in the case of the sun and earth.

This seems a satisfactory proof. But it might be objected that in this reasoning we assume the truth of the law of gravitation, which some may consider as an hypothesis and not a demonstrated truth. This objection would be reasonable. The proof I have given, although a very strong one, may appear to some not quite conclusive, so astronomers have sought for still stronger evidence, and this they have found in a remarkable phenomenon known as the "aberration of light." If the earth moves round the sun it may be easily proved, from the sun's known distance, that the velocity in its orbit must be about  $18\frac{1}{2}$  miles per second. The velocity at which light travels has been found by several methods, and is known to be about 186,000 miles per second. Now the light coming to us from a star

combined with the earth's motion causes an apparent displacement in the position of the star, as seen from the earth. This displacement is of course very small, owing to the comparatively slow motion of the earth, and can only be detected by very accurate instruments, but there is no doubt whatever that it exists. Were the earth at rest—as some argue—no such displacement could possibly occur. We therefore conclude that the earth is in motion; and as the observed "aberration" agrees exactly with the theoretical velocity, we conclude that the earth really does revolve round the sun. The supposed motion of the sun round the earth would *utterly* fail to explain the observed phenomenon,<sup>1</sup> and is therefore unworthy of serious consideration.

The planets also revolve round the sun, as we know from observation, and this is another argument in favour of the earth's motion, "the argument from analogy," as it is called. Why should the earth be the only body in the Solar System which remains at rest? This is a strong argument, but the argument from the "aberration of the stars" seems to me far stronger; indeed it is altogether unanswerable, and to the astronomer and mathematician the earth's motion round the sun is a *demonstrated fact*.

The earth's path round the sun is not a perfect circle. It is nearly so, but not exactly. The real

<sup>1</sup> This is the case with most of the hypotheses known as "paradoxes." They completely fail to explain observed phenomena, and are therefore worthless.

form is that of an ellipse, having what mathematicians call a small eccentricity. An ellipse is an oval figure with two points on its longer axis called the *foci*, and the sum of the distances of any point from the two foci is a constant quantity, and equal to the longer axis of the ellipse.

The point where the longer and shorter axis of the ellipse cross at right angles is called the *centre* of the ellipse. The distance from the centre to either focus is called the "eccentricity" of the ellipse. The greater the distance of the focus from the centre the more elongated is the ellipse. The orbits of comets are generally of this shape. The less the distance of the focus from the centre of the figure, the more nearly the ellipse approaches a circle in form; and when the focus coincides with the centre the ellipse becomes a circle. In fact the circle is only a particular case of the ellipse, or, as a mathematician would express it, the circle is the limiting form of the ellipse when the eccentricity is reduced to zero.

If we call the mean distance or semi-axis major 1, the present eccentricity will be expressed by the decimal fraction 0.0168, or very nearly by the vulgar fraction  $\frac{1}{60}$ . In other words, if we divide the semi-axis major into 60 equal parts, the distance of each focus from the centre will be equal to 1 of these parts. If therefore we assume that the earth's mean distance from the sun is 92,890,000 miles,<sup>1</sup> we find that the distance of the sun (which is placed in one of the

<sup>1</sup> Corresponding to a "solar parallax" of 8.80".




FIG. 7.—MOTION IN AN ELLIPTIC ORBIT.



foci) from the centre of the ellipse is (92,890,000 divided by 60, or) about 1,550,000 miles. Now if we subtract this from the mean distance, we obtain 91,340,000 miles from the earth's least distance from the sun, and if we add it, we find 94,440,000 miles for the earth's greatest distance from the sun. This difference, it will be seen, amounts to over 3 millions of miles, so that at one time of the year we are 3 millions of miles nearer to the sun than at the opposite point of the earth's orbit. The point of nearest approach is called the *perihelion*, and the point of greatest distance the *aphelion*. See diagram (fig. 7), which represents an ellipse of which the foci are S and F. Let S represent the focus in which the sun is situated. Then P will be the *perihelion*, and A the *aphelion*, CS the eccentricity, and  $SM=CP$  the mean distance. It is evident that SP, the perihelion distance, will be equal to  $CP-CS$ , and SA, the aphelion distance, equal to  $AC+CS$ , or  $CP+CS$ . The eccentricity of the ellipse shown in the diagram is considerably greater than that of the large planets, but is about the same as that of the minor planet. Polyhymnia (No. 33), the eccentricity of whose orbit is about 0.34 (or about one-third of the semi-axis major) as in the diagram.

It might be supposed that our nearest approach to the sun would occur in summer, but curious to say we are 3 millions of miles nearer the sun in December than in June.

This seems at first sight an extraordinary fact, but

in reality the difference of distance has little or nothing to do with the heat of the summer and the cold of winter. The difference of temperature is really due to the difference in the height of the sun above the horizon, and the length of the day. In the summer the days being very long, and the nights short, the earth and air are heated for a longer time than they are cooled during the night. The reverse of this takes place in winter.

The other planets also move in elliptical orbits round the sun, and all move in accordance with three laws which were discovered by the famous astronomer, Kepler, in the 17th century, and which are called "Kepler's Laws." The first of these laws is that which we have just considered, namely, that all the planets revolve round the sun in ellipses, the sun being situated in one of the foci. The second law is that the *radius vector*, or line joining the sun and planet, *describes equal areas in equal times*. The third law is, "*the squares of the periodic times are proportional to the cubes of the mean distances from the sun.*"

In the diagram (fig. 7) the *radius vector* is represented by SP, SM, or SA. The length of the *radius vector* depends upon the distance of the planet from the sun at any time. Now, the velocity at "perihelion" P is greatest, and that at "aphelion" A least. If in the diagram BP represents the arc traversed near perihelion in any given time, then near aphelion the distance described in the *same* time will be an

arc AD of such a length as to make the area ASD equal to the area PSB. This is true for every point in the orbit, the velocity varying so as to make the area swept over by the *radius vector* a constant quantity for equal times.

The time occupied by the planet in moving from E' through P to E will be less than in moving from E through A to E', as the area E'PE is less than the remainder of the ellipse. For this reason the winter in England is somewhat shorter than the summer.

To give an illustration of the third law of Kepler. Take the case of the earth and Mars. The periodic times are about 365 and 687 days, and the mean distances from the sun are in the proportion of 1 and 1.5237 respectively. Then we have the proportion—

$$365^2 : 687^2 :: 1^3 : (1.5237)^3$$

or 133,225 : 471,969 :: 1 : 3.5375 nearly.

Let us now consider the cause of the seasons. I explained in a former chapter that if a perpendicular be supposed drawn through the centre of the earth perpendicular to the plane of the ecliptic, then the earth's axis is inclined at an angle of  $23\frac{1}{2}$  degrees to this line, or at an angle of  $66\frac{1}{2}$  degrees to the plane of the orbit (the ecliptic). See fig. 6. To this inclination all the seasons of the year are due. If the axis of the earth were perpendicular to its orbit, it will be easily seen that the days and nights would be of equal length all over the earth, and at all times of the year. The earth's position with reference to the sun would then always remain the same, and




FIG. 8.—THE SEASONS.



there would then be no seasons. But as the earth's axis is inclined to the plane of its orbit, and always remains parallel to itself (always pointing to the celestial pole), it will be seen from the diagram (fig. 8), which is a perspective view of the earth's orbit round the sun, that at one season of the year the northern hemisphere is inclined *towards* the sun, and 6 months after is inclined *away* from the sun. In the southern hemisphere of course the same thing occurs, but in summer, when the North Pole is turned towards the sun, the South Pole is inclined away from it, and *vice versa*. Hence, when it is summer in England, it is winter in Australia and countries south of the equator, and their summer is our winter. At the intermediate positions C and D both hemispheres are similarly situated with reference to the sun, and at these times—March 21st and September 21st—day and night are of equal length all over the earth. These are called—for this reason—the *equinoxes*.

The curved lines on the earth's figure in the diagram represent the tropics. It will be seen that at A in the regions north of the northern tropic—called the “tropic of Cancer”—the greater portion of the diurnal path falls in the illuminated portion, and the smaller portion in the dark part. North of a certain point *p*, the latitude of which is evidently  $66\frac{1}{2}$  degrees, there is no portion in the dark part, and here there is no night in summer. It is “the Land of the Midnight Sun.” The reverse of this takes

place at B, where the whole region between *p* and the North Pole N is in the dark part, and then there is no day in those regions. The same thing occurs in the regions near the South Pole, but of course the seasons are there reversed, as at other places south of the equator.

The variation in the sun's distance will, however, have some slight effect on the temperature. It has this effect, that in the northern hemisphere the diminished distance of the sun in winter and the increased distance in summer mitigates, to some extent, the cold of winter and the heat of summer.

It will be seen from the diagram that in the southern hemisphere the summer occurs at the point of nearest approach to the sun or perihelion, and the winter occurs at aphelion. This will have the effect of *increasing* the summer temperature and diminishing the winter temperature, which is the reverse of what happens in the northern hemisphere.

Although the summer temperature of the southern hemisphere is greater than that of the northern, the total amount of solar heat received by each hemisphere during its summer season will be the same, for the following reason. When a planet, like the earth, revolves round the sun in an elliptical orbit, the velocity is greatest at the *perihelion* and least at the *aphelion*. Hence, as the English summer occurs at *aphelion*, when the earth is moving slowly, the summer will be somewhat longer than the winter. The reverse of this takes place in southern latitudes.

There the summer occurring at *perihelion*, the summer is *shorter* than the winter, and although the temperature shown by the thermometer is higher, it lasts for a shorter time, and the greater heat is counterbalanced by its shorter duration. At least this would be theoretically the case; but the effect is modified by other laws and local and geographical causes at different places.

But it may be asked, How is it known that the sun is nearer to the earth in the English winter than it is in summer?

This is very easily proved; for it is found that when the sun's apparent diameter is measured with astronomical instruments it is greater in winter than it is in summer, and the reader knows that the nearer an object is to the eye the larger it appears.

At the terrestrial equator there is but little change of seasons, for there the celestial equator passes through the zenith or point exactly overhead, so that on the 21st of June the sun is  $23\frac{1}{2}$  degrees north of the zenith, and on the 21st of December  $23\frac{1}{2}$  degrees south. The altitude, therefore, of the sun at noon is, on the earth's equator, never less than  $66\frac{1}{2}$  degrees, so that the inhabitants have a sort of perpetual summer. In fact, as the sun is on the celestial equator, or in the zenith twice a year, namely, on March 21st and September 21st, there are really *two* hot summers, with a milder summer between, and real winter is unknown.

## CHAPTER V.

### THE EQUATION OF TIME AND LEAP YEAR.

THE student will now understand the motions of the earth. The earth has a diurnal motion on its axis from west to east in about 24 hours, which produces the phenomena of day and night, and also the apparent daily motion of the sun, moon, and stars in the opposite direction, from east to west. In addition to this daily motion the earth has an annual motion round the sun, also from west to east, which produces the apparent motion of the sun among the fixed stars from west to east, and this motion, combined with the inclination of the earth's axis to the plane of its orbit, produces the changes of the seasons.

We will now proceed to consider the difference between *mean* time and *apparent* time. The earth is popularly supposed to rotate on its axis in 24 hours, but this is not exactly true. The correct period is 23h. 56m. 4s.

For the ordinary purposes of life it is found convenient to consider the day as the interval which elapses between two successive returns of the



sun to the meridian. If the sun were apparently fixed in the sky this interval would be exactly equal to the period of the earth's rotation, but, as you have learned, the sun has an *apparent* motion from west to east among the fixed stars. Hence, owing to this easterly motion, it will not come to the meridian exactly after one rotation of the earth, but a few minutes later. The solar day is therefore longer than the sidereal day (or period between two successive returns of the same star to the meridian). It is this solar day which is divided into 24 hours.

The length of the *solar* day is not, however, constant. It would be so if the sun moved along the celestial equator with a *uniform* velocity. But the sun does not move along the equator, but along the ecliptic, which is inclined to the equator, and its motion is not quite uniform, owing to the elliptic shape of the earth's orbit. In fact its apparent velocity is constantly varying to a small extent, being greatest in the English winter, and least in summer. The time shown by a sun dial is for this reason constantly varying.

The time shown by a sun dial is "apparent solar time," and a correction must always be made to the time found by such an instrument. This correction is called *the Equation of Time*. The result is called the "mean solar time."

This "equation of time" vanishes on four days of the year, namely, April 15th, June 15th, August 31st, and December 24th. I must explain, however, that for

astronomical purposes clocks are constructed to show "sidereal time," that is, the time which elapses between two successive returns of the same fixed star to the meridian. This period of 23h. 56m. 4s. is divided into 24 hours, so that a sidereal clock apparently "gains" nearly 4 minutes a day. But these clocks are only used in observatories.

I will now explain the necessity for a "leap year." The length of the year is not 365 days *exactly*, but about  $365\frac{1}{4}$  days, or, more exactly, 365 days, 6 hours, 9 minutes,  $9\frac{1}{2}$  seconds. This is called the "sidereal" year. The mean "solar" or "tropical" year is the interval which elapses between the passage of the sun through an equinoctial point and back to the same point. Were the equinoxes fixed points in the sky the "solar" and "sidereal" years would be of exactly the same length; but as these points have—owing to a motion called the "precession of the equinoxes"—a slow movement from east to west, it follows that the equinoctial point moves to meet the sun, and therefore the "solar" year is shorter than the "sidereal." The length of the solar year is 365 days, 5 hours, 48 minutes,  $46\frac{3}{4}$  seconds, or somewhat less than  $365\frac{1}{4}$  days. To allow for this quarter-day one day is added every four years. This year is called Leap Year, and contains 366 days, the other three having only 365 days each. Every year divisible by 4 is a leap year.

The year is, however, not *exactly*  $365\frac{1}{4}$  days, and this would cause some error in the course of time.

It would make a difference of about one whole day in 130 years, or about 3 days in 400 years. This is corrected by omitting a leap year at the beginning of each century, when the year—although divisible by 4—is not divisible by 400. Thus the years 1700, 1800, and 1900 are common years, but 2000 will be a leap year.

1900 will *not* be a leap year, although it can be divided by 4. 1896 will be a leap year, and then there will not be another till the year 1904.

This change in the method of reckoning was made in the year 1752 in this country, when the beginning of the year was also changed from March 25th to January 1st. It is called the "New Style." The "Old Style" ("O.S.") is still retained in Russia and Greece.

Aug  
100  
887  
1110  
1184  
1187  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225  
2226  
2227  
2228  
2229  
2230  
2231  
2232  
2233  
2234  
2235  
2236  
2237  
2238  
2239  
2240  
2241  
2242  
2243  
2244  
2245  
2246  
2247  
2248  
2249  
2250  
2251  
2252  
2253  
2254  
2255  
2256  
2257  
2258  
2259  
2260  
2261  
2262  
2263  
2264  
2265  
2266  
2267  
2268  
2269  
2270  
2271  
2272  
2273  
2274  
2275  
2276  
2277  
2278  
2279  
2280  
2281  
2282  
2283  
2284  
2285  
2286  
2287  
2288  
2289  
2290  
2291  
2292  
2293  
2294  
2295  
2296  
2297  
2298  
2299  
2300  
2301  
2302  
2303  
2304  
2305  
2306  
2307  
2308  
2309  
2310  
2311  
2312  
2313  
2314  
2315  
2316  
2317  
2318  
2319  
2320  
2321  
2322  
2323  
2324  
2325  
2326  
2327  
2328  
2329  
2330  
2331  
2332  
2333  
2334  
2335  
2336  
2337  
2338  
2339  
2340  
2341  
2342  
2343  
2344  
2345  
2346  
2347  
2348  
2349  
2350  
2351  
2352  
2353  
2354  
2355  
2356  
2357  
2358  
2359  
2360  
2361  
2362  
2363  
2364  
2365  
2366  
2367  
2368  
2369  
2370  
2371  
2372  
2373  
2374  
2375  
2376  
2377  
2378  
2379  
2380  
2381  
2382  
2383  
2384  
2385  
2386  
2387  
2388  
2389  
2390  
2391  
2392  
2393  
2394  
2395  
2396  
2397  
2398  
2399  
2400  
2401  
2402  
2403  
2404  
2405  
2406  
2407  
2408  
2409  
2410  
2411  
2412  
2413  
2414  
2415  
2416  
2417  
2418  
2419  
2420  
2421  
2422  
2423  
2424  
2425  
2426  
2427  
2428  
2429  
2430  
2431  
2432  
2433  
2434  
2435  
2436  
2437  
2438  
2439  
2440  
2441  
2442  
2443  
2444  
2445  
2446  
2447  
2448  
2449  
2450  
2451  
2452  
2453  
2454  
2455  
2456  
2457  
2458  
2459  
2460  
2461  
2462  
2463  
2464  
2465  
2466  
2467  
2468  
2469  
2470  
2471  
2472  
2473  
2474  
2475  
2476  
2477  
2478  
2479  
2480  
2481  
2482  
2483  
2484  
2485  
2486  
2487  
2488  
2489  
2490  
2491  
2492  
2493  
2494  
2495  
2496  
2497  
2498  
2499  
2500  
2501  
2502  
2503  
2504  
2505  
2506  
2507  
2508  
2509  
2510  
2511  
2512  
2513  
2514  
2515  
2516  
2517  
2518  
2519  
2520  
2521  
2522  
2523  
2524  
2525  
2526  
2527  
2528  
2529  
2530  
2531  
2532  
2533  
2534  
2535  
2536  
2537  
2538  
2539  
2540  
2541  
2542  
2543  
2544  
2545  
2546  
2547  
2548  
2549  
2550  
2551  
2552  
2553  
2554  
2555  
2556  
2557  
2558  
2559  
2560  
2561  
2562  
2563  
2564  
2565  
2566  
2567  
2568  
2569  
2570  
2571  
2572  
2573  
2574  
2575  
2576  
2577  
2578  
2579  
2580  
2581  
2582  
2583  
2584  
2585  
2586  
2587  
2588  
2589  
2590  
2591  
2592  
2593  
2594  
2595  
2596  
2597  
2598  
2599  
2600  
2601  
2602  
2603  
2604  
2605  
2606  
2607  
2608  
2609  
2610  
2611  
2612  
2613  
2614  
2615  
2616  
2617  
2618  
2619  
2620  
2621  
2622  
2623  
2624  
2625  
2626  
2627  
2628  
2629  
2630  
2631  
2632  
2633  
2634  
2635  
2636  
2637  
2638  
2639  
2640  
2641  
2642  
2643  
2644  
2645  
2646  
2647  
2648  
2649  
2650  
2651  
2652  
2653  
2654  
2655  
2656  
2657  
2658  
2659  
2660  
2661  
2662  
2663  
2664  
2665  
2666  
2667  
2668  
2669  
2670  
2671  
2672  
2673  
2674  
2675  
2676  
2677  
2678  
2679  
2680  
2681  
2682  
2683  
2684  
2685  
2686  
2687  
2688  
2689  
2690  
2691  
2692  
2693  
2694  
2695  
2696  
2697  
2698  
2699  
2700  
2701  
2702  
2703  
2704  
2705  
2706  
2707  
2708  
2709  
2710  
2711  
2712  
2713  
2714  
2715  
2716  
2717  
2718  
2719  
2720  
2721  
2722  
2723  
2724  
2725  
2726  
2727  
2728  
2729  
2730  
2731  
2732  
2733  
2734  
2735  
2736  
2737  
2738  
2739  
2740  
2741  
2742  
2743  
2744  
2745  
2746  
2747  
2748  
2749  
2750  
2751  
2752  
2753  
2754  
2755  
2756  
2757  
2758  
2759  
2760  
2761  
2762  
2763  
2764  
2765  
2766  
2767  
2768  
2769  
2770  
2771  
2772  
2773  
2774  
2775  
2776  
2777  
2778  
2779  
2780  
2781  
2782  
2783  
2784  
2785  
2786  
2787  
2788  
2789  
2790  
2791  
2792  
2793  
2794  
2795  
2796  
2797  
2798  
2799  
2800  
2801  
2802  
2803  
2804  
2805  
2806  
2807  
2808  
2809  
2810  
2811  
2812  
2813  
2814  
2815  
2816  
2817  
2818  
2819  
2820  
2821  
2822  
2823  
2824  
2825  
2826  
2827  
2828  
2829  
2830  
2831  
2832  
2833  
2834  
2835  
2836  
2837  
2838  
2839  
2840  
2841  
2842  
2843  
2844  
2845  
2846  
2847  
2848  
2849  
2850  
2851  
2852  
2853  
2854  
2855  
2856  
2857  
2858  
2859  
2860  
2861  
2862  
2863  
2864  
2865  
2866  
2867  
2868  
2869  
2870  
2871  
2872  
2873  
2874  
2875  
2876  
2877  
2878  
2879  
2880  
2881  
2882  
2883  
2884  
2885  
2886  
2887  
2888  
2889  
2890  
2891  
2892  
2893  
2894  
2895  
2896  
2897  
2898  
2899  
2900  
2901  
2902  
2903  
2904  
2905  
2906  
2907  
2908  
2909  
2910  
2911  
2912  
2913  
2914  
2915  
2916  
2917  
2918  
2919  
2920  
2921  
2922  
2923  
2924  
2925  
2926  
2927  
2928  
2929  
2930  
2931  
2932  
2933  
2934  
2935  
2936  
2937  
2938  
2939  
2940  
2941  
2942  
2943  
2944  
2945  
2946  
2947  
2948  
2949  
2950  
2951  
2952  
2953  
2954  
2955  
2956  
2957  
2958  
2959  
2960  
2961  
2962  
2963  
2964  
2965  
2966  
2967  
2968  
2969  
2970  
2971  
2972  
2973  
2974  
2975  
2976  
2977  
2978  
2979  
2980  
2981  
2982  
2983  
2984  
2985  
2986  
2987  
2988  
2989  
2990  
2991  
2992  
2993  
2994  
2995  
2996  
2997  
2998  
2999  
3000  
3001  
3002  
3003  
3004  
3005  
3006  
3007  
3008  
3009  
3010  
3011  
3012  
3013  
3014  
3015  
3016  
3017  
3018  
3019  
3020  
3021  
3022  
3023  
3024  
3025  
3026  
3027  
3028  
3029  
3030  
3031  
3032  
3033  
3034  
3035  
3036  
3037  
3038  
3039  
3040  
3041  
3042  
3043  
3044  
3045  
3046  
3047  
3048  
3049  
3050  
3051  
3052  
3053  
3054  
3055  
3056  
3057  
3058  
3059  
3060  
3061  
3062  
3063  
3064  
3065  
3066  
3067  
3068  
3069  
3070  
3071  
3072  
3073  
3074  
3075  
3076  
3077  
3078  
3079  
3080  
3081  
3082  
3083  
3084  
3085  
3086  
3087  
3088  
3089  
3090  
3091  
3092  
3093  
3094  
3095  
3096  
3097  
3098  
3099  
3100  
3101  
3102  
3103  
3104  
3105  
3106  
3107  
3108  
3109  
3110  
3111  
3112  
3113  
3114  
3115  
3116  
3117  
3118  
3119  
3120  
3121  
3122  
3123  
3124  
3125  
3126  
3127  
3128  
3129  
3130  
3131  
3132  
3133  
3134  
3135  
3136  
3137  
3138  
3139  
3140  
3141  
3142  
3143  
3144  
3145  
3146  
3147  
3148  
3149  
3150  
3151  
3152  
3153  
3154  
3155  
3156  
3157  
3158  
3159  
3160  
3161  
3162  
3163  
3164  
3165  
3166  
3167  
3168  
3169  
3170  
3171  
3172  
3173  
3174  
3175  
3176



It is more than 700 times the mass of all the planets and satellites put together.

The sun rotates on its axis like the earth. But the period of rotation is much longer—about  $25\frac{1}{4}$  days. This is ascertained by watching the motion of spots on its surface.

The exact nature of these spots has not been well ascertained. Numerous theories have been advanced to account for these curious phenomena; but they are believed to be openings in the outer envelope of the sun, and are known to be most prevalent on the sun's surface every 11 years, and less numerous during intermediate years.

It was formerly believed that the solar light and heat were due to some form of combustion, but the enormous quantity of fuel necessary to support this combustion, and the great difficulty of explaining where this vast amount of fuel came from, has led astronomers to abandon the theory. How, then, is the expenditure of light and heat maintained? It is now almost generally admitted that the heat and light are due to the contraction or shrinkage of the solar mass. It may be shown mathematically that this shrinkage is quite adequate to produce the observed effects, and it certainly seems to be the most satisfactory theory which has yet been advanced. The theory is due to the eminent German physicist, Helmholtz, and has been ably supported by Sir William Thomson.

But, it may be asked, if shrinkage can produce

light and heat, to what cause is this shrinkage due? The answer to this is simple. The shrinkage is due to the immense attraction of the sun's great mass on its component particles, and owing to the comparatively small density of the sun this attraction is able to act more effectively on the sun's mass than it could do on the heavier materials which form our earth.

But, it may be said, if the sun is shrinking in this way, it must be gradually diminishing in size. Could not this diminution be detected by astronomical instruments? We might naturally fancy it could, but Sir W. Thomson, who ably advocates Helmholtz's theory, has shown that the shrinkage necessary to produce the observed amount of light and heat is so small that the diminution in the sun's apparent diameter, even in the long period of 2000 years, could scarcely be detected by our most accurate instruments. And of course we have no measurements of the sun's diameter dating nearly so far back as 2000 years.

This theory seems to be a simple and satisfactory one, and as we know that the enormous gravitating power of the sun *must* produce shrinkage in its mass, and that the shrinkage of a large mass *must* produce light and heat, there seems to be no necessity to look further for an explanation of the "mystery of the sun." In fact "the fuel of the sun" is no longer a mystery, and the solar light and heat find a satisfactory explanation in the "gravitation

theory." Of course the *cause* of gravitation itself still remains as great a mystery as ever.

Although the sun is *comparatively* close to us when compared with the immense interval which separates us from even the nearest of the fixed stars, still its distance is *absolutely* very great. Light with its enormous velocity of 186,000 miles per second, takes over 8 minutes to reach us from the sun. A cannon ball, moving with a velocity of 1,500 feet per second, would take over 10 years to traverse the distance, and a railway train, travelling at the speed of 60 miles an hour, would take about 176 years in its journey across the vast space which separates us from the solar orb.

## CHAPTER VII.

### THE MOON.

NEXT to the sun the moon is the most conspicuous object in the heavens, and being our nearest celestial neighbour forms of course an object of especial interest. Its diameter is about 2,160 miles, and its mean distance from the earth about 238,800 miles.

In addition to being an interesting and beautiful object, I have explained in a previous chapter that it is useful for finding the longitude at sea, as well as for giving light at night.

The division of time into months evidently had its origin in the period of the moon's revolution round the earth. This period is not exactly a calendar month, which is an arbitrary division of time. The period in which the moon moves from any point in the sky round to the same point again is 27 days, 7 hours, 43 minutes. This is called the *sidereal* revolution, but while the moon is performing its revolution round the earth, the earth itself is proceeding on its annual journey round the sun. For this reason the moon has to travel some








FIG. 10.—PHASES OF THE MOON.

moon. In the diagram (fig. 10) let E represent the earth, S the sun, and suppose the moon to revolve in the direction of the arrows. Then neglecting the earth's motion round the sun (which will not affect the explanation); when the moon is at N, between the earth and S, its illuminated side is turned towards the sun, and its dark side towards the earth. In this position it becomes invisible, and is called "new moon." When it advances to the point A, a portion of the illuminated side becomes visible in the form of a crescent, and when it arrives at the point B, half the moon is visible. It is then said to be in *quadrature* (because the angle between the sun and moon is then a right angle) or in popular language the "First Quarter." When it reaches the point C, more than half the illuminated side comes into view, and it is then said to be *gibbous*. At F, directly opposite to the sun, or in *opposition*, we see the full disc, and it is then called "Full Moon." Still proceeding in its course these changes are repeated in inverse order, being "gibbous" at D, in "quadrature" at E, called "Last Quarter"; and in the crescent shape again at G, a few days before "New Moon." After "new moon," as it appears to the left (or east) of the sun at A, the crescent is seen in the evening after sunset; and at G before "new moon," being to the right (or west) of the sun, the crescent is seen in the morning before sunrise.

But it may be asked, How is it that we only



see one side of the moon? This is due to the fact that the moon rotates on its axis in the same time that she revolves round the earth (see fig. 11). But it is not strictly true that we never see any portion of the opposite hemisphere. Owing to the slightly elliptical shape of the moon's orbit her motion round the earth is not quite uniform, while her rotation on her axis is perfectly so. This gives rise to a phenomenon called *libration*, which sometimes brings a portion of the further hemisphere into view, near the east and west limbs.<sup>1</sup>

If we represent the whole spherical surface by 1, the area of the portion which always remains invisible will be 0.4111, or about 41 per cent. Or in other words, if the whole surface be represented by 100, then 41 will represent the part permanently invisible, and 59 the portion known to observers.<sup>2</sup>

When the moon is a crescent the student may have often seen the dark side, when the air is clear. This is simply due to sunshine reflected from the earth, or "earthshine," as it is sometimes called. The earth is very much larger than the moon, and consequently, as seen from the moon, the earth would appear very much larger than the moon appears to us. In fact it would appear about

<sup>1</sup> There is another small libration which brings portions near the poles into view.

<sup>2</sup> Taking the moon's diameter at 2,160 miles, we have the total area of its surface, 14,657,407 square miles, and hence the area which remains permanently invisible will be 6,025,660 square miles.




FIG. 11.—ROTATION OF THE MOON ON ITS AXIS.

13 times as large, and therefore gives considerably more light to the moon than the moon does to us.

The earth as seen from the moon would show similar phases; with this difference, that when the moon is "full" to us the earth is "new" to the moon, and at "new" moon the earth is "full." Now it is clear why the dark side is faintly visible a little after "new moon," for the earth is then nearly "full" to the moon, and lights up its surface much more brilliantly than full moonshine on the earth.

The existence of any forms of life on the moon's surface is very improbable. Observations show that the moon is devoid of air and water, which are absolutely necessary to most forms of life. If it has an attenuated atmosphere, it is so rarefied that its existence cannot be detected by our most delicate observations.

The phenomenon of the "harvest moon" is familiar to most people. This is best explained by reference to a celestial globe, but I will try and make the matter clear in another way. The "harvest moon" is the full moon which occurs on or *nearest* to the 21st of September (the autumnal equinox), and is so called because it is useful at that season in the evenings for getting in the crops. The moon being then in the sign Aries,<sup>1</sup> rising due east, and the

<sup>1</sup> Owing to the "precession of the equinoxes" the "sign" Aries is now in the constellation Pisces, and the sign Libra in Virgo.



FIGS. 12, 13.—TO ILLUSTRATE PHENOMENON OF HARVEST MOON.



sun entering Libra, and setting due west, the southern half of the ecliptic is then above the horizon, and the northern half below it, and the ecliptic then makes the smallest possible angle with the horizon. Now the moon moving nearly in the ecliptic its rising is less retarded for several days than at other times. Perhaps a diagram will make the matter clearer. In fig. 12, let A, B, C, D represent the position of the moon on four consecutive days at the time of harvest moon (FEB the horizon), and in fig. 13, let G, H, I, K represent its position on four consecutive days, when its apparent path in the heavens makes a larger angle with the horizon MLH. Then it is evident that the spaces through which it must rise at C and D, namely CE and DF, are much less than the corresponding spaces IL and KM in the second figure. The distances AB, BC, CD are equal, and each equal to GH, HI, and IK. Hence the time of rising in fig. 12 is much less delayed than in fig. 13.

At the equator, as the north and south poles are on the horizon, the ecliptic makes the same angle with the horizon at the vernal and autumnal equinoxes, so the phenomenon of the harvest moon is not apparent. Indeed, at the equator the harvest moon is not required. The further north we go the more marked is the phenomenon.<sup>1</sup>

The student may have noticed that in winter the

<sup>1</sup> Of course, within the arctic and antarctic circles the moon remains constantly above the horizon during the winter.

moon is high in the heavens, and in summer is low down near the southern horizon. This is due to its path lying near the ecliptic, which is high in winter and low in summer. In winter when its light is most required it is high in the sky, and in summer when the bright nights render moonlight unnecessary it is low. In this will clearly be seen the evidence of Divine wisdom.

## CHAPTER VIII.

### ECLIPSES.

ECLIPSES of the sun and moon are very interesting phenomena, and are of tolerably frequent occurrence (especially those of the moon). I will begin with those of the sun, as the nature of these is very simple. An eclipse of the sun is caused by the moon passing between the sun and earth.

This evidently can only occur at "new moon," when the moon is between the sun and earth. But it may be asked, How is it that an eclipse of the sun does not happen at every full moon? This is due to the fact that the moon's orbit is inclined to the ecliptic at a small angle (about  $5\frac{1}{4}$  degrees). For this reason the moon generally passes either a little above or a little below the sun at the time of new moon.

Under what circumstances, then, can an eclipse of the sun occur? The moon's orbit intersects the ecliptic at two points which are called the *nodes*. When the new moon occurs in or near one of these nodes an eclipse of the sun takes place.

### ECLIPSES.

73

The apparent diameter of the moon is not always the same as that of the sun. Owing to the elliptical shape of the orbits of the earth and moon, the moon's apparent diameter is sometimes greater and sometimes less than that of the sun. When the moon is at her greatest distance from the sun, and her apparent diameter consequently at its smallest, if it happens to be *in* the node at the time of new moon we have the phenomenon known as an "annular eclipse," a narrow ring of sunlight being visible all round the moon's black disc. When the moon is nearest to the earth and in the node, we have a *total* eclipse of the sun—a phenomenon of very rare occurrence, at least in Great Britain. In all other cases the eclipse is only a *partial* one.

A total eclipse cannot last long. Under the most favourable conditions the duration of the total phase cannot much exceed 7 minutes, and it is generally considerably less than this.

The last total eclipse visible in England took place on May 22, 1724, and the next will not happen till August 11, 1999,<sup>1</sup> when the sun will be totally eclipsed for about two minutes in the south-east of England, at Plymouth, Torquay, Weymouth, &c. An "annular" eclipse will be visible from the Shetland and Orkney Islands on April 8, 1921.<sup>1</sup>

<sup>1</sup> See "Eclipses and Transits in Future Years," by the Rev. S. J. Johnson, M.A., F.R.A.S. (Parker and Co., Southampton Street, Strand). The solar eclipse of June 29, 1927, will be total for a few seconds in the north of England.



The next solar eclipse of any magnitude visible in England will occur on May 28, 1900, when seven-tenths of the sun's disc will be eclipsed in the south of England.

A total eclipse of the sun is one of the grandest and most imposing sights in Nature. During the total phase a ring of light with rays is seen round the moon's disc. This is called the solar "corona," and is a glowing appendage surrounding the sun, which, owing to the intense glare of sunlight, is not visible on ordinary occasions.

An eclipse of the moon is caused by the moon passing through the earth's shadow, and can of course only occur at "full moon," when the sun, earth, and moon are in the same straight line. This does not happen at every "full moon," for the same reason that a solar eclipse does not occur at every "new moon," namely, the inclination of the moon's orbit to the plane of the ecliptic, which enables it to pass sometimes above and sometimes below the shadow of the earth.

A lunar eclipse lasts much longer than a solar one.

During an eclipse of the moon a lunar spectator would see the sun eclipsed by the earth's disc, which appears about 13 times larger than the moon does to us. It can therefore hide the sun for a much longer time. Another way of explaining the matter is that the diameter of the earth's shadow at the moon's distance is very considerable, so that the moon takes a long time to pass through it. For a




FIG. 14.—ECLIPSE OF THE MOON.

sphere, when the source of light is a globe larger than the sphere (as in the case of the sun and earth) the shape of the shadow is always a cone. If the sun and earth were of the same size, the shadow would be a cylinder of infinite length, but as the sun is much the larger, the earth's shadow is a cone, as will be seen by the accompanying diagram (fig. 14, which is of course not drawn to scale, but merely intended to represent the general principle of a lunar eclipse). S represents the sun, E the earth, and M the moon. The darkly shaded triangle represents the conical shadow, which is called the *umbra*, and the lighter shade is called the *penumbra*. Within the conical shadow the whole of the sun's disc is hidden from the moon, but in the *penumbra* a portion of the sun's disc only is eclipsed, as will be seen by the lines touching the edges of the sun and earth. Thus, if from any point P, within the penumbra, a line PC, parallel to AD, be drawn, touching the earth's surface and meeting the sun's surface at C, then to a spectator at P only the portion AC will be hidden by the earth, while the portion CB will remain uneclipsed. Hence there will be only a *partial* eclipse at P; and similarly for the other points within the limits of the *penumbra*.

From any given point on the earth's surface eclipses of the moon are more frequently *seen* than those of the sun; because when the moon is either partially or totally eclipsed, the eclipse is visible

from every point on the earth's surface from which the moon is visible at the time, but eclipses of the sun are only visible at certain points. In fact, an eclipse of the moon is a *real* eclipse, that is, a real darkening of the lunar surface, whereas those of the sun are merely occultations by the moon, and may be total in one place, and quite invisible at others. In reality, however, eclipses of the sun are more numerous than those of the moon. Out of 70 eclipses which usually occur in about 18 years, the average number of eclipses of the sun is 41, and of lunar eclipses 29.

The period of 18 years forms what is termed a "cycle," which was known to the ancient Greeks under the name of the *Saros*. During this period eclipses recur in nearly the same order.

As I have already explained, eclipses take place near the *nodes* of the moon's orbit, or the points at which the lunar orbit intersects the plane of the ecliptic. Now it is found from observation that these nodes do not maintain a fixed position, but have a rapid *retrograde* motion on the ecliptic<sup>1</sup> in a period of about  $18\frac{1}{2}$  years.<sup>2</sup> The apparent motion of the node is therefore in the *opposite* direction to that of the sun among the fixed stars. Consequently the sun, in its apparent annual motion through the heavens, will meet the same node in less than a

<sup>1</sup> Or contrary to the order of the signs of the Zodiac.

<sup>2</sup> The exact period of revolution of the node is 6793.391 days.



year, or in about 346.62 days. The moon's synodical revolution (or the period from one new or full moon to the next) is 29 days, 12 hours, 44 minutes. Now 223 of these periods amount to about 6,585 $\frac{1}{3}$  days, while 19 periods of 346.62 days amount to 6,585 $\frac{2}{3}$  days. Hence it follows that at the end of this period, or about 18 years, 11 days, the sun, the moon, and the moon's node will return nearly to the same position with reference to each other, and consequently if an eclipse occurred at the beginning of the period, a similar, or nearly similar, eclipse will recur at the end.

Another cycle known to the ancient astronomers is termed the *metonic cycle*. This consists of 19 years of 365 $\frac{1}{4}$  days, or 6,939 $\frac{3}{4}$  days, which is very nearly equal to 235 synodical revolutions of the moon.

## CHAPTER IX.

### MERCURY.

HAVING now described the earth and moon, we will proceed to consider the other planets and satellites of the Solar System, commencing with Mercury, the nearest to the sun. Mercury and Venus, the two planets which revolve round the sun in orbits included *within* the earth's orbit, are termed "inferior planets."

It is known that these planets revolve *inside* the earth's orbit, from the very simple reason that they are found by observation to be constantly changing their places among the stars, and are never seen further from the sun than a limited distance. For example, if a man were to walk round a tree, placed in the centre of a circular enclosure, and a person standing at the boundary of the enclosure observed that the pedestrian was never seen in the opposite direction to that of the tree, he would conclude that his circular path round the tree was *included* within the boundary of the enclosure. For exactly the same reason the planets Mercury and Venus are

never seen in the part of the heavens opposite to the sun's place.

The greatest distance that Mercury can be seen from the sun, or the "greatest elongation"—as it is termed—of the planet cannot exceed  $27\frac{1}{4}$  degrees, or less than one-third of the distance from the horizon to the zenith, and it rarely exceeds 18 degrees (or one-fifth of the same distance). For this reason the planet is seldom seen with the naked eye, being only visible in the morning and evening twilight.

The mean distance of Mercury from the sun is about 36 millions of miles, and the period of revolution round the sun—or the length of its year—is about 88 of our days. The orbit is, however, very elliptical, and for this reason the distance varies from about 27 millions to 43 millions of miles. Its orbit is inclined to the ecliptic at the angle of about 7 degrees.

Bessel supposed the period of rotation on its axis to be about 24 hours, but this is very uncertain. Professor Schiaparelli has recently announced his opinion, based on 10 years' observation, that the rotation is similar to that of the moon—that is, that the period of rotation on its axis is equal to the period of revolution round the sun, so that the planet always turns the same side towards the sun. This result, if confirmed, is certainly very interesting and unexpected.

The diameter of Mercury is about 3,000 miles, so that in volume it is the smallest of all the large

primary planets.<sup>1</sup> But in density it is very heavy—about 7 times the weight of water (that of the earth being about  $5\frac{1}{2}$ ).

Owing to the intense heat of the sun we cannot suppose it to be inhabited by any forms of life, except perhaps close to the poles of the planet; or, if Schiaparelli's theory be correct, near the boundary line dividing the light from the dark side.

Mercury, seen with a telescope, shows phases similar to those of the moon, which proves that it shines—like the moon—by light reflected from the sun. Its apparent motion among the fixed stars resembles that of Venus, and this I will explain when speaking of that planet.

Mercury occasionally passes across the sun's disc like a small black spot. The last transit of Mercury occurred in November, 1881, and the next will take place on May 10, 1891, and November 10, 1894. These transits can of course only occur when the planet is in that portion of its orbit which lies between the earth and the sun, or when the planet is in *inferior* conjunction, as it is termed. When the planet passes behind the sun it is said to be in *superior* conjunction.

The student may ask, How is it that a transit does not occur at *every* inferior conjunction? For the same reason that an eclipse of the sun does not take place at every new moon, namely, that the orbit of

<sup>1</sup> All the "minor planets," revolving between Mars and Jupiter, are, however, much smaller.



Mercury is inclined to the ecliptic, and consequently the planet sometimes passes a little above the sun, and sometimes a little below it, when in conjunction. A transit can only occur when the planet is in or near its node (or inclination of its orbit plane with the plane of the ecliptic), and this rarely occurs.

## CHAPTER X.

### VENUS.

NEXT to Mercury comes the brilliant planet Venus, which is often seen in the evening sky after sunset. It is the brightest of all the planets, occasionally so bright as to cast a shadow on a white surface!

It is considerably larger than Mercury, and but little smaller than the earth, its diameter being about 7,700 miles (the earth's mean diameter being 7,912). It revolves round the sun at a mean distance of 67 millions of miles, in a period of about 225 days. Its period of rotation on its axis is supposed to be about  $23\frac{1}{4}$  hours, but this is doubtful,<sup>1</sup> owing to the difficulty of seeing any well-defined markings on its surface. The position of the axis of rotation is still more uncertain.

The "greatest elongation" of Venus from the sun can never exceed about 47 degrees, or about half the distance from the horizon to the zenith.

<sup>1</sup> Schiaparelli is of opinion that Venus—like Mercury—rotates on its axis in the same time that it revolves round the sun.

Venus sometimes appears brighter than at others. This is due to its varying distance from the earth. When it is near "inferior conjunction," or nearly between the earth and sun, its distance from the earth is only 26 millions of miles,<sup>1</sup> whereas when in "superior conjunction," or on the other side of the sun, its distance is about 160 millions of miles,<sup>2</sup> or its greatest distance is about 6 times its least distance. When near its least distance, however, it shows a thin crescent as the moon does when near the sun, whereas when near its greatest distance it exhibits a full face, so that its brilliancy is not so greatly diminished in the latter case as its increased distance in this position might at first sight lead us to expect.

I said in the last chapter that I would explain the apparent motion of Mercury and Venus among the fixed stars. At first sight the apparent motion seems somewhat complicated, but it really admits of a simple explanation, as will be seen by studying the accompanying diagram (fig. 15), which is carefully drawn to scale (supposing the orbits to be circular, as they nearly are). A, B, C, D, &c., represent successive positions of the earth at intervals of 10 days, and A' B', C', D' the corresponding positions of Venus. The outer circle is supposed to represent the star sphere, and *a, b, c, d, &c.*, are the successive positions of the planet as seen from the earth at A, B, C, D, &c., respectively; S being the position of the sun, and the arrows representing the direction of motion.

<sup>1</sup> 67 deducted from 93.

<sup>2</sup> 67 added to 93.

A' is the point of "greatest elongation" as seen from the earth's position at A, the angle AA'S being a right angle (and AA' a tangent to the orbit of Venus at A'). As the small letters *a, b, c, d, &c.*, representing the apparent positions of Venus among the stars, are in alphabetical order, an examination of the diagram will show that the apparent motion of the planet is *direct* from *a* to *e* (or in the order of the signs). At some point between *e* and *f* the direct motion ceases, and the planet becomes, for a short time, stationary,<sup>1</sup> and afterwards retrograde, so that when the earth arrives at F (and Venus at F') the planet has retrograded to *f*. This retrograde motion continues through inferior conjunction (HH'h) up to a point near L, where the planet again becomes stationary, and the motion, shortly after, again *direct*, and this direct motion is maintained through "superior conjunction." From A to H Venus is an evening star, and from I to N a morning star.<sup>2</sup> Its varying distance from the earth will be seen from the diagram, AA' being considerably greater than HH', where the planet is nearly at its minimum distance.

<sup>1</sup> In some text books on astronomy it is stated that a planet is *stationary* when it is moving nearly in a line towards an observer on the earth. This is, however, quite incorrect. At the point A in the diagram Venus is moving directly towards the earth, but it is not then stationary; on the contrary, it has at this point a *rapid direct motion*.

<sup>2</sup> Of course the planet would have been an evening star for a considerable time before it reached the point A', and remains a morning star for a long time after it has passed the point N'.



The student may have heard of transits of Venus being employed to find the sun's distance from the earth. The actual details of the calculation are very complicated, but the general principle of the method is sufficiently simple, and will be easily understood. In the diagram (fig. 15) illustrating the relative motions of the earth and Venus, it will be seen that when the planet is at its greatest elongation at  $A'$ , if we measure the angle of elongation  $SAA'$ , which can be easily done, we can (since the angle at  $A'$  is a right angle) find, by the rules of plain trigonometry, the ratio of  $SA'$  to  $SA$ . Now  $SA'$  is the distance of Venus from the sun, and  $SA$  is the earth's distance from the sun. The ratio of these distances can therefore be found by observation and simple calculation.<sup>1</sup> (It could also be determined by Kepler's third law.)

When Venus is projected on the sun's disc during a transit, its apparent position as seen from stations widely separated on the earth's surface may be ascertained by careful observation, and the observed displacement enables us to calculate the sun's distance from the earth, as will be seen from the accompanying diagram (fig. 16). Let  $A$  be a station in the earth's northern hemisphere, and  $B$  a station in the southern hemisphere, separated by a diameter of the earth.

<sup>1</sup>  $\frac{SA'}{SA} = \sin SAA'$ . Hence if we make  $SA = 1$ , we have  $SA' = \sin SAA'$ . The relative distance of Venus and the earth being about 67 and 93 we have  $\frac{67}{93} = 0.7204 = \sin SAA'$ , and therefore  $SAA' = 46^\circ 5'$  approximately.




FIG. 15.—RELATIVE POSITIONS OF THE EARTH AND VENUS FROM GREATEST "ELONGATION" OF VENUS, THROUGH INFERIOR CONJUNCTION.

Then seen from A the centre of Venus, C, will be projected on the sun's disc at D, and seen from B it will appear at E.<sup>1</sup> Hence the total apparent displacement is ED. Now the alternate angles at C are (by Euclid) equal, and therefore ED is to AB as CD is to AC. But the latter ratio is about 67 to 26. Hence ED is about  $2\frac{1}{2}$  times AB, and therefore the angle subtended by ED at the earth is about  $2\frac{1}{2}$  times the angle subtended by AB at the sun. From this, knowing AB, the sun's distance from the earth can be found.

The distance ED may be found by noting the exact times at which Venus, moving in its orbit in the direction of the arrow, enters the sun's disc at F and H ("ingress") and leaves it at G and K ("egress"). From these intervals of time the lengths of the lines FG and HK may be found, and hence the value of ED may be computed.

This process is, in theory, simple enough, but in practice several corrections must be made before the necessary accuracy is attainable. For instance, the exact motions of Venus and the earth in their orbits during the transit, the rotation of the earth on its axis, &c., must be taken into account, and these details make the calculations complicated and troublesome.

Transits of Venus only occur after long intervals,

<sup>1</sup> The apparent displacement is of course greatly exaggerated, as the relative sizes of the earth, Venus, and the sun are not drawn to scale.



FIG. 16.—TRANSIT OF VENUS.



and then usually in pairs. These intervals are 8, 122, 8, 105, 8, 122 years, &c. The last pair occurred on Dec. 8, 1874, and Dec. 6, 1882, and the next two will not take place till June 7, 2004, and June 5, 2012. I had the good fortune to see both the transits in 1874 and 1882 myself. The former I observed in India, and the latter in the West of Ireland. Both were seen in a clear sky.<sup>1</sup>

<sup>1</sup> "Planetary and Stellar Studies," p. 32.

## CHAPTER XI.

### MARS.

NEXT to Mercury and Venus, in order of distance from the sun, comes our own earth, but as we have already discussed its dimensions and motions, we will now pass on to the "red planet" Mars, which is the next planet outside the earth. It revolves round the sun in a period of about 687 days, at a mean distance of about 141 millions of miles. The shape of its orbit is, however, more elliptical than that of the earth (see fig. 17). For this reason it is sometimes much nearer to the earth than at others, and consequently varies considerably in brilliancy at different oppositions.

Mars is larger than Mercury, but considerably smaller than the earth and Venus. Its diameter is about 4,200 miles, so that in volume the earth is about  $6\frac{1}{2}$  times larger.

It rotates on its axis, like the earth and the other planets, and the period of rotation, or the length of

its day, has been determined with great accuracy from observations of markings on its surface. The latest and most reliable computation makes this period 24 hours, 37 minutes, 22.66 seconds, so that the length of the day in Mars is known to within a tenth of a single second!

Mars is sometimes stationary among the stars, and sometimes retrograde, as in the case of Venus. This will be understood from the accompanying diagram (fig. 17). Starting from A, when Mars at A' is seen in the sky at *a*, forming a right angle with the sun at S (or in "quadrature," as it is termed), the relative positions of the earth and Mars are shown at intervals of 10 days (as in the diagram of Venus). It will be seen that the lines joining the positions of the earth and Mars seem to *radiate* outwards from A to I, at which point they become nearly parallel, and afterwards apparently *converge* to the point O, when they again become parallel, and then again radiate outwards to the point X (when Mars is again in quadrature). Now, as parallel lines drawn from any two points in the earth's orbit to meet the celestial sphere point to the same star (owing to the immense distance of the stars), the radiation of the lines AA'a, BB'b, CC'c, &c., *outwards* shows that in these positions Mars has a *direct* motion among the stars. When the lines become parallel, near I, the planet becomes for a short time *stationary*, and when they converge, between J and N, the planet has an apparent *retrograde* motion. After passing OO', the lines PP'p,




FIG. 17.—RELATIVE POSITIONS OF THE EARTH AND MARS BEFORE AND AFTER "OPPOSITION" OF MARS.



QQ'q, RR'r again radiate outwards, and the planet's motion again becomes *direct*.<sup>1</sup>

Near the "quadratures" ( $a$  and  $x$  in the diagram) Mars is not seen with a full face—owing to the relative positions of the planet and the earth. At these points the planet appears somewhat "gibbous" in shape, like the moon a few days before and after full moon.

I may here explain the meaning of the terms *geocentric* and *heliocentric* place of a planet. The *geocentric* place is the apparent position as seen from the earth (or rather the earth's centre), and the "*heliocentric*" place its position as seen from the sun (or rather the sun's centre). For example, in fig. 17 the geocentric place of Mars at C' is  $c$ , but its heliocentric place is  $s$ . At the point L' the "geocentric" and "heliocentric" places are the same. When the planet is in "opposition," the sun, earth, and planet are then in the same straight line, or nearly so, and then the "geocentric" and "heliocentric" places coincide.

I spoke of "markings" on the disc of Mars. These markings are considered—and with great probability—to indicate the existence of land and

<sup>1</sup> In some text books on astronomy it is erroneously stated that a superior planet (like Mars) will appear *stationary* when the earth is advancing directly towards the planet. An inspection of fig. 17 will show that at the point A the earth is advancing *directly* towards Mars, and at X *directly* away from it. But at these points the planet is *not* stationary, but in *direct* motion. At the real stationary points, near I and O, the earth is *not* moving straight towards the planet.

water on the surface of the planet. Elaborate maps have been made of Mars on this supposition, showing various continents and oceans. If this view be correct—as it probably is—the areas of land and water on the planet are about equal. This is a different arrangement to that of the earth. On the earth the water surface is about 3 times as great as that of the land.

For many years Mars was supposed to have no moons, and Tennyson speaks of "the snowy poles of moonless Mars." Two very small satellites were, however, discovered in 1877 by Prof. Asaph Hall with the 26-inch refractor of the Washington Observatory. These are supposed to be about 6 and 7 miles in diameter. The inner moon has been called Phobos, and the outer one Deimos. Phobos is distant from the centre of Mars only 6,000 miles, and revolves round its primary in the unusually short period of 7 hours, 39 minutes. The distance of Deimos is about 15,000 miles, and its period of revolution about 30 hours, 18 minutes. These minute moons are among the smallest members of the Solar System.

## CHAPTER XII.

## THE MINOR PLANETS.

BETWEEN Mars and Jupiter a number of very small planets revolve round the sun. The orbits of many of these are very elliptical in shape, much more so than those of the larger planets. The first of these interesting discoveries was Ceres, which was found by Piazzi at Palermo on the first day of the present century, Jan. 1, 1801. In 1802 Olbers discovered Pallas. In 1804 Juno was found by Harding, and in 1807 Vesta by Olbers. No more were detected till 1845, when Astræa was discovered by Hencke, and in 1847 Hebe, by the same observer. Since that time yearly additions have been made to the list, and up to the present (October, 1890) no less than 299 have been found.

Vesta, the brightest of all, has been occasionally seen with the unassisted vision. Ceres may be seen with a good opera-glass, when favourably situated; but the great majority are only visible with a telescope.

The eminent astronomer Stone thinks that the

diameter of the first 71 minor planets varies from 60 to 214 miles. The fainter ones are of course probably still smaller, and Le Verrier has shown that the mass of all combined cannot exceed one-fourth of the earth's mass.

The orbits of some of them are inclined at a considerable angle to the plane of ecliptic. For instance, the orbit of Pallas has an inclination of nearly 35 degrees, those of Euphrosyne and Istria about  $26\frac{1}{2}$  degrees, Gallia nearly 26 degrees, and Æthra about 25 degrees. Others, however, have only a small inclination. Their orbits also intersect each other in a curious way, very different from those of the larger planets, which — when even at their nearest — are separated by a considerable distance.



## CHAPTER XIII.

### JUPITER.

WE next come to the "giant planet" Jupiter, the largest member of the Solar System, exceeding in volume all the other planets put together.

Jupiter is much larger than Venus, but owing to his much greater distance from the earth and sun he appears fainter. His mean diameter is about 87,000 miles, so that in volume he exceeds the earth about 1,373 times, and Venus about 1,490 times. In density, however, Jupiter is much lighter than either the earth or Venus, its specific gravity (water equal 1) being only 1.30, that of the earth being 5.67, and Venus about 4.82.

The mean distance of Jupiter from the sun is about 483 millions of miles, and it revolves round the solar orb in a period of 11 years and 10 months.

Jupiter has a very rapid rotation on its axis for so large a globe, the period of rotation being about 9 hours, 55 minutes, 37 seconds. This gives a velocity at the planet's equator of about 8 miles a second! This velocity is about 28 times greater than the

speed of rotation at the earth's equator. This rapid rotation has been shown mathematically to explain the observed compression at the north and south poles of Jupiter's disc, which is very evident in a good telescope.

This compression is much greater than in the case of the earth, which is, as I said in a former chapter, nearly a sphere. The polar diameter of Jupiter is shorter than the equatorial diameter in the proportion of about 100 to 106½.

Jupiter is attended by four satellites or moons, which revolve round the planet in the same way that the planets revolve round the sun. One of them—the second in order of distance from the planet—is almost exactly the same size as the moon, its diameter being about 2,100 miles. The others are larger, No. I. being about 2,400 miles in diameter, No. III. 3,400, or somewhat larger than Mercury, and No. IV. about 2,900 miles. The following are their approximate distances from Jupiter, and periods of revolution round their primary :—

Satellite.	Distance from Jupiter.	Period.
	MILES.	D. H. M.
I.	274,000	1 18 39
II.	437,000	3 13 14
III.	697,000	7 3 43
IV.	1,226,000	16 16 32

The first satellite is therefore somewhat farther from Jupiter than the moon is from the earth, and the others at considerably greater distances. Like

our moon, these satellites apparently rotate on their axes in the same time that they revolve round their primary. The length of the day on Satellite I. is therefore  $42\frac{1}{2}$  hours, and on the others longer.

These satellites are subject to eclipse in Jupiter's shadow in the same way that our moon is eclipsed in the shadow of the earth; but, unlike those of our moon, these eclipses are of frequent occurrence as seen from Jupiter. The reason of this is that the satellites revolve round Jupiter approximately in the plane of the planet's equator, which nearly coincides with the plane of its orbit (or "ecliptic," as we call it in the case of the earth).

The three inner satellites are eclipsed at every revolution, but the outer one, IV., occasionally escapes being eclipsed owing to the fact that its orbit is slightly inclined to those of the other satellites. A remarkable relation exists between the motions of the three inner satellites, from which it follows that all three can never be eclipsed together.

These eclipses are visible from the earth with the aid of a telescope, and the accompanying diagram (fig. 18) will illustrate the matter. S represents the sun; P the planet throwing a dark shadow behind it; A, B, C, D, E, F, G the earth's orbit, and *a*, *b*, *c*, *d* the orbit of one of the satellites, the arrows showing the direction of motion. Now when the earth is at A, and the satellite at *a*, it will enter the planet's shadow (called the *immersion*) and be eclipsed, as seen from the earth. When it arrives at *b* it will emerge




FIG. 18.—ECLIPSES OF JUPITER'S SATELLITES, AS SEEN FROM THE EARTH.



from the shadow (called the *emersion*). When it reaches *c* it will pass behind the planet's disc, as seen from A. This is termed an *occultation*; and when it arrives at *d* the satellite will reappear on the opposite side of the planet. In the position shown in the diagram Jupiter is seen to the right of the sun, and is then a morning star. The eclipses and occultations will then occur, as shown in the figure, or the eclipses first and the occultations afterwards. When the earth advances to G the planet will be an evening star, and the above phenomena will take place in reverse order, namely, the occultations first and the eclipses afterwards. Near D, when Jupiter is in "opposition" (rising at sunset), the occultations only will be visible, the eclipses occurring when the satellite is behind the planet. In the case of the first and second satellites, if the *immersion* can be seen, the *emersion* is invisible, and *vice versa*. Another interesting phenomenon is the transit of a satellite across the disc of Jupiter. This occurs when the satellite is passing from *e* to *f*, as seen from A. In these transits the satellites are sometimes seen as black spots on Jupiter's disc. This seems curious, as they must be—in that position—fully illuminated by the sun's light. An attempt has been made to explain the fact by supposing that the planet possesses some inherent light of its own. A better explanation, however, seems to be that the apparent blackness of the satellites in transit is due to the difference in light reflecting power of Jupiter and the satellites—an effect of contrast in fact.

The surface of Jupiter is, however, very bright, not much inferior to that of white paper, and from *this* fact it seems very probable that the planet shines partly by inherent light—in other words, that its surface is red hot. It cannot, therefore, be inhabited, but its satellites may possibly form the abodes of some forms of life.

Most people have heard of Jupiter's "belts." These are darkish bands across its surface, as seen in a telescope, usually lying nearly parallel to the planet's equator, and generally of various shades of yellow and brown. They are not, however, constant features, their details being occasionally subject to rapid changes. Some astronomers consider that these "belts" represent the body of the planet seen through openings in its cloudy envelope.

## CHAPTER XIV.

## SATURN.

OUTSIDE Jupiter's orbit revolves the "ringed planet" Saturn, which, until the discovery of Uranus, was considered the outermost planet of the Solar System.

Its distance from the sun is much greater than that of Jupiter. Its mean distance is about 886 millions of miles, or about  $9\frac{1}{2}$  times the earth's mean distance from the sun. Its period of revolution is about  $29\frac{1}{2}$  years. As heat and light vary *inversely* as the square of the distance, we have the intensity of the heat and light received by the earth equal to 9.5 multiplied by 9.5, or about 91 times as great as Saturn experiences.

Saturn is a very large planet, but not quite so large as Jupiter, its mean diameter being about 72,000 miles, or over 9 times greater than that of the earth. As, however, the volumes of spheres vary as the *cubes* of their diameter it follows that Saturn exceeds the earth in volume about 797 times. In density, however, it is very light, its specific gravity being only 0.68 (water equal 1), that of the earth being 5.67. For this reason its mass (or quantity of matter) is

only 95 times greater than that of the earth. It has therefore little more than half the density of Jupiter. In fact it is the lightest planet of the Solar System, as far as we know at present.

Saturn is attended by eight moons, which have been named Mimas, Enceladus, Tethys, Dione, Rhea, Titan, Hyperion, and Japetus, in order of distance from the planet, Mimas being the nearest. Their distances from the centre of Saturn range from about 122,700 miles to about 2,350,000 miles, and their periods of revolution round their primary from about  $22\frac{1}{2}$  hours to 79 days,  $7\frac{3}{4}$  hours.

Titan, the largest, is supposed to be about 3,300 miles in diameter, or a little larger than Mercury. The diameters of the others are somewhat doubtful, but the outer one, Japetus, is next in order of size. The two interior ones, Mimas and Enceladus, are very faint objects, and can only be seen in large telescopes.

According to Prof. Asaph Hall, Saturn rotates on its axis in 10 hours, 14 minutes, 24 seconds. It is, like Jupiter, considerably flattened at the poles, the ratio of the equatorial to the polar diameter being about as 12 to 11.

The rings of Saturn form the most unique and interesting feature in this wonderful system. They consist of two flat luminous rings, with a dark division between them, and enclosing a semi-luminous ring, known as the "dusky" or "crape" ring. These completely surround Saturn, but do not anywhere touch the planet. From some measures made in the years



1880-81 by Dr. Meyer at the Geneva Observatory I find that the exterior diameter of the outer ring is about 173,500 miles; interior diameter of the middle ring, 112,440 miles; interior diameter of dusky ring, 90,810 miles; width between dusky ring and ball of planet, 7,632 miles. The width of the dark division between the bright rings, called Cassini's division—after its discoverer—is about 1,700 miles.

An eclipse of the satellite Japetus in the shadows of the globe, crape ring, and bright ring of Saturn, took place on Nov. 1, 1889. This interesting and rare phenomenon was well observed by Mr. E. E. Barnard at the Lick Observatory. His observations show that "the crape ring is truly transparent—the sunlight sifting through it. The particles composing it cut off an appreciable quantity of sunlight. They cluster more thickly, or the crape ring is denser as it approaches the bright rings. . . . So far as the penetration of the solar rays is concerned, the bright ring is fully as opaque as the globe of *Saturn* itself."<sup>1</sup>

It may be proved, however, from mechanical principles, that the rings cannot possibly be solid. The hypothesis that they may be liquid is also untenable. The theory now generally adopted by astronomers is that they consist of a swarm of minute satellites which revolve round the planet in separate orbits.

Comparatively speaking the rings are very thin—probably not more than 100 miles in thickness. When the plane of the rings passes through the earth,

<sup>1</sup> *Monthly Notices*, R.A.S., January, 1890.

which occurs at intervals of  $14\frac{2}{3}$  years (half the period of revolution), the edge only is visible, and in this position the ring system, being so thin, disappears, even in a telescope of considerable power.

Saturn is probably in a red-hot state, like Jupiter. The light reflecting power of the planet is considerably less than that of Jupiter, but judging from its small density, and the appearance of "belts" on its surface, similar to those of Jupiter, it seems probable that the planet is in a heated condition. *Most* of Saturn's light is, however, merely reflected sunlight.

## CHAPTER XV.

## URANUS AND NEPTUNE.

WE now come to a planet which was unknown to the ancient astronomers. Uranus was discovered by Sir W. Herschel on March 13, 1781, and was at first mistaken for a comet, but its planetary nature was soon determined, and it was found to revolve round the sun in an orbit exterior to that of Saturn.

Its mean distance from the sun is about 1,782 millions of miles, or more than double the distance of Saturn. It takes about 84 years to perform one revolution round the sun.

Uranus is very much larger than the earth, but considerably smaller than Jupiter and Saturn, its diameter being about 33,600 miles. In density, however, Uranus is much lighter than the earth, its specific gravity being about equal to that of water.

Uranus most probably rotates on its axis, like the other planets, but owing to its small disc, and the difficulty of seeing any markings on its surface, even with large telescopes, the period has not been ascertained. Judging, however, from the analogy of Jupiter

and Saturn, it seems probable that the period does not differ much from 10 hours.

There are four known satellites which have been named Ariel, Umbriel, Titania, and Oberon, in order of their distance from the planet. The distances of these moons range from about 125,000 miles to 382,000 miles. Unlike the satellites of the planets already considered, the planes of their orbits are nearly at right angles to the plane of the ecliptic, and the motion is retrograde—that is, they move from east to west, instead of from west to east.

When favourably situated Uranus shines as a star of about  $5\frac{1}{2}$  magnitude, and is just visible to the naked eye. It may be seen, however, with an opera-glass on any clear night, if its position is accurately known. At present (Oct., 1890) it is situated a little to the east of the bright star Spica ( *$\alpha$  Virginis*).

From recent observations with the spectroscope, it seems probable that—like Jupiter and Saturn—Uranus is still in a highly heated condition. From its observed brightness I find that its brilliancy is considerably greater than theory would lead us to expect in a planet so far from the sun. Before its discovery as a planet, Uranus was observed no less than 20 times, and recorded as a fixed star on each occasion:

NEPTUNE.—Next to Uranus comes Neptune, the most distant known member of the planetary system. It was discovered on Sept. 23, 1846; but its existence had previously been predicted, and its position pointed



out by two young mathematicians, Adams in England, and Le Verrier in France. After the discovery of Uranus, irregularities were observed in its motion which the theory of gravitation could not account for. The discrepancies between the computed and observed places of Uranus at last became so considerable that astronomers were led to suspect the existence of an unknown planet outside Uranus, which, by its attraction, was disturbing the motion of Herschel's planet. The enormously difficult mathematical problem of calculating the position of this hypothetical planet was independently undertaken by the two mathematicians I have just named. The result was the telescopic discovery of the planet by Galle at Berlin close to the computed position. This was a splendid scientific triumph, and in the opinion of some astronomers the discovery of Neptune forms the most brilliant achievement in astronomy since the discovery of the law of gravitation by Sir Isaac Newton.

The mean distance of Neptune is about 2,792 millions of miles, or about 30 times the earth's distance from the sun. The period of revolution is about  $164\frac{1}{2}$  years, or nearly double that of Uranus.

Neptune is somewhat larger than Uranus, its diameter being nearly 38,000 miles. In density, however, it is rather lighter, being about 0.9 (water equal 1). The period of revolution on its axis is unknown.

Only one satellite of Neptune has, up to the present,



FIG. 19.—POSITION OF NEPTUNE OBSERVED BY LALANDE MAY 8TH AND 10TH, 1795.

been discovered. Like the Uranian satellites its motion is retrograde, and the plane of its orbit is inclined to the plane of the planet's orbit at a considerable angle. It revolves round Neptune in 5 days, 21 hours, at a distance of about 220,000 miles, or somewhat less than that of our moon. The fact of its being visible at all at the great distance which separates us from Neptune makes it probable that it is a body of considerable size.

As in the case of Uranus, Neptune was—previous to its discovery as a planet—observed and recorded on several occasions as a fixed star. The accompanying diagram (fig. 19) shows its position as observed by Lalande on May 10, 1795.

## CHAPTER XVI.

### COMETS.

THE student will now like to hear something about the comets. Some of them form members of the Solar System, but the very bright and large ones are, as a rule, only seen once, and must be considered as merely occasional visitors to our system.

The comets which belong to the Solar System travel in elliptic orbits; but these ellipses are generally very elongated—that is, one of the axes of the ellipse—the *major axis*, as it is called—is very much longer than the other, or, as a mathematician would express it, the eccentricity of the ellipse is very great. Other comets, however, are supposed to travel in a curve called the parabola—that is, a curve which does not return into itself, but has two similar branches which run out to infinity. In this case a comet having once passed round the sun would go off into space never to return. As, however, it is difficult to determine whether the small portion of the orbit near the sun belongs to an ellipse or parabola—both curves nearly coinciding near the focus—it is not always certain whether a



comet is really travelling in a parabolic orbit. The orbit may, in many cases, be really an ellipse of great length.

Among the comets known to be regular *members* of the Solar System, there are two groups, one consisting of small comets with periods ranging from  $3\frac{1}{2}$  years (Encke's) to  $13\frac{3}{4}$  years (Tuttle's). These can only be observed with a telescope. The second group consists of 5 comets having periods of  $67\frac{3}{4}$  to  $76\frac{3}{4}$  years. The most remarkable of these is that known as Halley's, which has a period of about 76 years. Apparitions of this comet have been traced back to B.C. 11. Its last appearance took place in 1835, and its next return is due in the year 1910.

The periods of some of the large comets have also been calculated, but the great length of the computed periods renders them very uncertain. The famous comet of 1811 was calculated by Argelander to have a period of over 3,000 years! The magnificent comet of 1858, known as Donati's, has been computed to have a period of about 2,000 years. The great comet of 1861 seems to have been moving in an elliptic orbit, with a period of about 419 years. Coggia's comet of 1874 is supposed to have a period of over 10,000 years! That of 1882 was well observed, and its period has been fixed at 772 years, with considerable approach to accuracy.

Comets have been for ages a mystery to astronomers; but it is now thought possible that the *head* may consist of a cloud of meteoric stones, and that

the *tail* is due to electrical repulsion existing in the sun which drives out into space the minuter particles of matter contained in the nucleus. This theory explains the observed fact that the tail is gradually formed as the comet approaches the sun, and afterwards disappears as the comet recedes into space.

Some comets approach the sun very closely; the great comets of 1680, 1843, and 1882 nearly grazed the surface of the sun. Those of 1843 and 1882 were seen with the naked eye close to the sun in broad daylight.

## CHAPTER XVII.

### DOUBLE AND BINARY STARS.

A *DOUBLE* star is one which appears to the naked eye as an ordinary single star, but when examined with a good telescope is seen to be composed of two stars very close together.

The number of these interesting objects now known to astronomers probably exceed 10,000. These, however, include objects of varying character; some being easily seen with small telescopes, while others cannot be clearly separated by the highest powers of the largest telescopes yet constructed!

There are some stars which may be seen double with the naked eye or an opera-glass, but these cannot properly be termed "double" stars, at least astronomers would not consider them so. Some observers speak of those visible to the unassisted vision as "naked eye doubles," but this is of course only a popular way of speaking. You may have noticed a small star close to the centre star in the tail

of the Great Bear (or handle of the "Plough"). This is called by some "Jack on the Middle Horse," and, though now clearly visible to good eyesight, it was called by the ancient astronomers Alcor, or the "test," as it was then considered rather a difficult object, and a test of keen vision. It may possibly have increased in brightness, but the evidence in favour of variation in its light at present does not seem to me conclusive.

Only a few stars are, however, close enough to look—to the naked eye—like real double stars, but I may mention the brightest star in the constellation Capricornus, which can be easily seen double with good eyesight on a clear night. The star known as  $\alpha$  Cygni (a little to the west of  $\alpha$  Cygni) is another example. The most southern of the stars in the "sword" of Orion,  $\iota$ , may also be seen double on fine nights, but this is perhaps a more difficult object than the others. The most severe test, however, visible in these latitudes is the star  $\epsilon$  Lyræ (near Vega), which some eyes can see double. It is, however, easily visible in a good opera-glass. This is a very remarkable star, as with a good telescope each of the components is seen to be again double, forming a "double-double" or quadruple star, and between the pairs are several fainter stars.

In many cases—especially in the wide pairs—the components are, in all probability, only apparently close together, and when one of the stars is very much fainter than the other it possibly lies at a vast distance



behind the brighter star.<sup>1</sup> There are many cases, however, in which the components are really close together, and therefore at nearly the same distance from the earth. In these double stars Sir William Herschel discovered that one of the components actually revolves round the other, in the same way that the earth and other planets revolve round the sun, and these are called Binary, or revolving double stars.

These are most interesting objects—revolving suns. Their periods of revolution vary from about 11 years to over 1,000 years. The shorter periods are, however, comparable with those of the larger planets, Jupiter, Saturn, Uranus, and Neptune.

I said that the orbits of the earth and the larger planets were ellipses, but nearly approaching the circular form. In the majority of the binary stars the orbit is a lengthened ellipse; some of them, indeed, resembling more the orbits of comets than planets. In others, however, the orbit does not differ more from a circle than that of Mercury, and in a few instances the orbit approaches even still closer to the circular form.

A considerable number of stars are known to be binary, and others are suspected, but the orbits of

<sup>1</sup> There are, however, notable exceptions to this rule; for instance, Sirius,  $\delta$  Cygni, 85 Pegasi, 99 Herculis, all of which form binary systems. I have quite recently computed an orbit for 99 Herculis and find a period of  $53\frac{1}{2}$  years, with an ellipse of large eccentricity.

only about 60 have been hitherto calculated by astronomers. Some of these are bright stars; others faint. The following are bright stars:—Sirius,  $\alpha$  Centauri, Castor,  $\gamma$  Leonis,  $\gamma$  Virginis,  $\zeta$  Sagittarii,  $\zeta$  Herculis,  $\beta$  Delphini,  $\delta$  Cygni,  $\eta$  Cassiopeiæ, and others.

The periods being so long these binary stars require to be watched for many years before any apparent change becomes visible. In some cases we have observations of the more remarkable, extending more than 100 years back.


In watching the motion of these revolving stars we do not see the real orbit described. From the principles of mechanics the student will understand that the component stars really revolve round their common centre of gravity, which lies somewhere between them. But, for convenience of measurement and calculation, one of the stars—the brighter of the two—is supposed to be at rest, and the whole of the observed motion to take place in the other. The orbit thus observed is called the *apparent* orbit, and even this does not represent the real motion, for as a rule the plane in which the components revolve is *inclined* to the line of sight, and only in the case where this plane happens to be at right angles to the line of sight—a rare case—do we ever see the *real* orbit of a binary star.

The *apparent* orbit observed differs considerably in shape from the real orbit. But both orbits are ellipses, or rather are assumed to be so, on the supposition

that the law of gravitation holds good in these systems as well as in our own Solar System, which is of course highly probable. According to Kepler's second law, equal areas are described in equal times. This is true for *both* the *apparent* and the *real* ellipse. In fact the law is true for a body moving in *any* curve round a centre of force.

The reader might like to see the apparent orbit of one of these binary stars. Here is a diagram (see fig. 20) of the apparent orbit of the companion of Sirius which I have myself recently computed.<sup>1</sup> It shows the form and position of the apparent ellipse, and the portion of the orbit which has been described by the small star since its discovery in 1862. Sirius is the brightest star in the heavens. It has a considerable "proper motion," and from irregularities in this motion, which could not otherwise be explained, astronomers came to the conclusion that its motion was disturbed by the attraction of some unknown companion. Shortly after this opinion was published a faint companion was discovered by the celebrated optician, Alvan Clark, in America; and careful observations since that time fully confirm the suspicion that it is actually revolving in an orbit round Sirius. In the diagram, C is the centre of the apparent ellipse, A the position of the bright star. The dates show the position of the faint star in the years given. If C and A be joined, and produced both ways, this line will represent the projection of the *major*, or longer

<sup>1</sup> *Monthly Notices*, Royal Astronomical Society, June, 1889.





axis, of the *real* ellipse, and P is the *periastron*, or point of nearest approach in the *real* orbit. It will be seen that the point marked *m* is the point of nearest approach of the components in the *apparent* orbit (M that of greatest distance), and does not coincide with the periastron point in the real orbit.<sup>1</sup>

When the real orbit is at right angles to the line of sight, the minimum distance is then at the periastron, and in one or two other cases; but as a general rule the point of nearest approach is *not* at the *periastron*.

The line marked in the diagram "line of nodes" is the line of intersection of the plane of the *real* orbit with the plane on which the orbit is projected, or the background of the sky (or the plane of the paper in the diagram). One half the orbit is *below* this plane, the other half *above* it; but it is impossible to decide from the observations of position which half is below and which above.

The orbits formerly computed for Sirius gave a period of 49 or 50 years, but I find that recent measures require a somewhat longer period—about  $58\frac{1}{2}$  years.

There are many other interesting binary stars. One of the most remarkable is  $\gamma$  Virginis, which has a period of about 185 years. During last century it

<sup>1</sup> The mistaken idea that the least distance *always* occurs at the *periastron* seems a not uncommon one. Some years since a well-known astronomer expressed his opinion that the famous binary star  $\alpha$  Centauri had not then reached the periastron, *because* the components had not yet reached their minimum distance!

was an easy object, even in small telescopes, but it gradually "closed up," and in the year 1836 became so close that the most powerful telescopes of that day could not show it as anything but a single star! It then "opened out" again, and is at present an easy object, even with a small telescope. The ellipse in this case is very elongated, very much indeed resembling the orbit of a comet. Another elongated orbit is that of the southern star  $\alpha$  Centauri—the nearest "fixed star" to the earth—as far as has yet been determined. The period is about 77 years. Another interesting binary is 70 Ophiuchi, for which I have also computed an orbit, and find a period of about 88 years.<sup>1</sup> The German astronomer, Schur, some years since found a period of about 94 years, but this does not satisfy recent observations. The bright star Castor also consists of two components, of which the period is about 1,000 years.

The shortest period known is  $\delta$  Equulei, for which the Russian computer, Wrublewsky, finds a period of about  $11\frac{1}{2}$  years. It is a very close pair, and can only be measured with the largest telescopes. Other binary stars with short periods are  $\zeta$  Sagittarii,  $\beta$  Delphini, 85 Pegasi, and 42 Comæ.

The calculation of these double star orbits is generally difficult and troublesome. The measures of the same star by different astronomers often differ so considerably that it requires much experience and patience to obtain anything like a satisfactory result.

<sup>1</sup> *Monthly Notices*, R.A.S., March, 1888.

In most cases also the portion of the ellipse hitherto described is so small that the exact ellipse, of which it forms a segment, must be uncertain.

There are some cases, however, in which a complete revolution has been performed. The star known as  $\xi$  Ursæ Majoris has completed a full revolution since its discovery by Sir W. Herschel in 1780. In 42 Comæ more than two complete revolutions have been performed; the period is about  $25\frac{3}{4}$  years. The components of  $\zeta$  Herculis have revolved nearly three times since 1782—period about 34 years. The orbits of these stars have consequently been ascertained with considerable accuracy.

## CHAPTER XVIII.

### VARIABLE STARS.

THE variable stars are those which are not constant in their light, but which vary in brightness. Some of them vary to a very great extent, but others only slightly. They form a very interesting and mysterious class of objects.

Over 200 stars are known to be certainly variable, and many others have been suspected.

Some of these variables may be observed with the naked eye. Perhaps the most remarkable of them all is that known as Mira, or the "wonderful" star, in Cetus. This extraordinary object has a period, from maximum to maximum, of about 331 days. When at its brightest it shines as a star of about the 2nd magnitude.<sup>1</sup> At the minima it diminishes to about the 9th magnitude.<sup>2</sup> The exact position of the star will be seen from the following

<sup>1</sup> Sometimes, however, it does not rise above the 4th magnitude.

<sup>2</sup> It always remains visible in a telescope of 3 inches aperture.



diagram (fig. 21), which will enable the student to find it in the sky without much difficulty.

Another remarkable object of this class is the star known as Algol, or  $\beta$  Persei. The fluctuations in the light of this star are very curious. The period is very short—about 2 days, 20 hours, 48 minutes, 51 seconds. For the greater portion of this period the light of the star remains steady at a little below the 2nd magnitude. It then begins to diminish in brightness, and in about 4 hours, 23 minutes is reduced to a star of about magnitude  $3\frac{3}{4}$ . At this brightness it remains for about 15 minutes, when the light again begins to increase, and in about  $5\frac{1}{4}$  hours it recovers its ordinary brilliancy. The accompanying diagram (fig. 22) will enable the student to easily find this curious star. As, however, the minima of light occur at all hours of the day and night, it is not an easy matter to "catch" it at its faintest. From the end of March to the beginning of August the star is not favourably situated for observation, being low down on the northern horizon during the early portion of the night. The epochs of minimum are given in Whittaker's Almanac.

The "Algol stars" form a very rare type of variable, only 9 having hitherto been detected. Most of these are small, and not easily observed, but the following are visible to the naked eye:  $\lambda$  Tauri and  $\delta$  Libræ. There is another interesting class of variables known as "short period" variables. These are in a constant state of variation in periods ranging



FIG. 21.—THE VICINITY OF MIRA ( $\theta$ ) CETI.

from 1 day to 17 days. Of these the following are tolerably bright stars:  $\zeta$  Geminorum, period about 10 days,  $3\frac{3}{4}$  hours;  $\beta$  Lyrae, period about 12.9 days;  $\eta$  Aquilæ, period 7 days, 4 hours, 14 minutes; and  $\delta$  Cephei, period 5 days, 8 hours, 47 minutes, 40 seconds.<sup>1</sup> Other smaller objects of this class are known, some of them very interesting, but the aid of an opera-glass is required to follow their fluctuations.

No very satisfactory theory has yet been advanced to account for the changes of brightness in stars of long period, but in the case of stars of the Algol type it is thought probable that the diminution of light is due to the interposition of a dark eclipsing satellite which, at the star's minimum, cuts off a portion of its light, and some recent spectroscopic observations by Vogel seem to show that this is really the solution of the mystery.

<sup>1</sup> Further particulars respecting these and other variable stars will be found in my "Scenery of the Heavens."



FIG. 22.—THE VICINITY OF ALGOL ( $\beta$  PERSEI).



## CHAPTER XIX

## THE NEBULÆ.

IN addition to the thousands of stars visible in the heavens, there are a large number of objects known as *nebulae*, from their hazy appearance. Over 8,000 are now known to astronomers, and doubtless there are many more.

A few of the nebulae are visible to the naked eye; for instance, the great nebulae in Andromeda and Orion; but the great majority are only visible with a good telescope.

Many of them, although usually classed under the general head of *nebulae*, are really clusters of small stars; others, or the true nebulae, are known—from observations of their light with the spectroscope—to consist of nothing but glowing gas, or mixture of gases, of which hydrogen is certainly one constituent.

The great nebula in the "Sword" of Orion is a mass of luminous gas. It may be well seen with a small telescope, but of course large instruments are necessary to show the details of this wonderful object. It has been recently photographed with great success,

No telescope has yet succeeded in resolving the great nebula in Andromedæ into stars, but the spectroscope shows that it is *not* gaseous. Probably it consists of very small stars, or possibly a swarm of meteorites so small as to be individually invisible at the immense distance which separates it from the earth. A wonderful photograph of this object recently taken by Mr. Roberts at Liverpool shows a number of rings partially separated from the central mass.

Several of the clusters may be seen with a small telescope. One in the constellation Gemini, a little north of the star  $\eta$  in that constellation, may be well seen with a 3-inch telescope, although, of course, larger instruments are necessary to show the fainter stars. The cluster known as 34 Messier, between  $\beta$  Persei (Algol) and 60 Andromedæ, is a very good object for a small telescope. I have seen stars sparkling in this cluster with a binocular field-glass. The double cluster near  $\chi$  Persei is another splendid object in a small instrument. This has been recently photographed at the Paris Observatory. Among the globular clusters, one between  $\eta$  and  $\xi$  Herculis is a magnificent object in a good telescope. This has also been photographed. Further particulars respecting the nebulae and clusters, and photographs of some of them, will be found in my "Scenery of the Heavens."

## INDEX.

- ADAMS, 110  
 Æthra, minor planet, 97  
 Alcor, 117  
 Aldebaran, 29  
 Algol, 126  
 Altair, 29  
 Andromeda, Great nebula in, 131  
 Annual motion of earth, 40  
 Annular eclipse, 73  
 Aphelion, 46  
 Apparent time, 53  
 Apparent orbit of a binary star, 120  
 Arcturus, 29, 30  
 Ariel, satellite of Uranus, 109  
 Asteroids, *see* MINOR PLANETS,  
 Astrea, minor planet, 96  
 Astronomy, Meaning of, 15  
 Aurige,  $\alpha$  (Capella), 29
- BALLOON, 23  
 Barnard, 106  
 Bayer, 29  
 Behrmann, 25  
 Bessel, 80  
 Binary stars, 118
- CANOPUS, 29  
 Capella, 29, 30  
 Capricornus,  $\alpha^1$  and  $\alpha^2$ , 117  
 Cassini, 106  
 "Cassiopeia, Chair of," 32
- Castor, 123  
 Catalogues of stars, 25  
 Centauri,  $\alpha$ , 29  
 Cephei,  $\delta$  (variable star), 128  
 Ceres, minor planet, 96  
 Ceti, Mira (variable star), 125  
 Clusters, Globular, 131  
 Coggia comet, 114  
 Coma Berenices, 42 (binary), 124  
 Comets, Nucleus of, 114  
     " Tail of, 115  
     " Remarkable, 114, 115  
 Corona, Solar, 74  
 "Crape ring" of Saturn, 105
- DAY and night of equal length, 50  
 Declination, 36  
 Deimos, satellite of Mars, 95  
 Delphini,  $\beta$  (binary star), 123  
 Diameter of earth, 19  
     " Jupiter, 98  
     " Mars, 91  
     " Mercury, 80  
     " Moon, 61  
     " Neptune, 110  
     " Saturn, 104  
     " Uranus, 108  
 Dione, satellite of Saturn, 105  
 Diurnal motion of earth, 20, 32  
     " " sun and stars, 20,



Division, Cassini's, in Saturn's ring, 106

Donati, Comet of, 114

"Dusky ring" of, 105

EARTH, diameter of, 19

" distance from sun, 44

" a globe, 16, 17

" Orbit of, 44

" Proofs of rotundity of, 16

" seen from moon, 74

" Specific gravity of, 98

Eccentricity of ellipse, 44

Eclipses of Jupiter's satellites, 100

" moon, 74

" sun, 72, 73

Ecliptic, 32

Ellipse, 44

Enceladus, satellite of Saturn, 105

Encke, Comet of, 114

Equation of time, 53, 54

Equator, 34

Equuleii,  $\delta$  (binary), 123

Ether of space, 28

Euphrosyne, minor planet, 97

"FLAT earth" theory untenable, 16

Figure of earth not flat, but globular, 16

Fixed stars, 25

Foucault's experiment, 21

GALLE, 110

Gallia, minor planet, 97

Gaseous nebulae, 130

Geminorum,  $\alpha$  (Castor), 30

"  $\beta$  (Pollux), 30

Globular clusters, 131

HALL, Prof., 95

Halley, Comet of, 114

Harvest moon, 68

Heat of sun, 58

" Jupiter, 103

" Saturn, 104

Hebe, minor planet, 96

Heis, 25, 26

Helmholtz, theory of sun's heat, 58,

59

Hemisphere, The sky a, 23, 24

Hencke, 96

Hercules, Globular cluster in, 131

Herculis, 99 (binary star), 118

Herschel, Sir W., 108, 124

Hévelius, 25

Hipparchus, 25

Horizon sensible, 22, 23

" rational, 22, 23

"Hull down," ship, 16, *Frontispiece*

Hyperion, satellite of Saturn, 105

INFERIOR PLANETS, 79

Infinity of space, 27, 28

JAPETUS, satellite of Saturn, 105, 106

Juno, minor planet, 96

Jupiter, "Belts" of, 103

" brightness of disc, 103

" Density of, 98

" Diameter of, 98

" Distance from sun, 98

" Period of, 98

" Probable heat of, 103

" Rotation of, 98

" Satellites of, 99

" Volume of, 98

KEPLER'S LAWS, 47, 48

LATITUDE, 34, 38

Leap Year, 55

Leonis,  $\alpha$  (Regulus), 32

Le Verrier, 110

Libration of moon, 66

Lick telescope, 27

Limited number of stars, 26, 27

Longitude, 39

"Lunar distances," 39

Lyra;  $\alpha$  (Vega), 29

"  $\beta$  (variable star), 128

"  $\epsilon$ , test for keen vision, 117

MAGNITUDES of stars, 28

Mars, Diameter of, 91

" Distance from sun, 91

" Period of, 91

" Rotation of, 91, 92

" Satellites of, 95

Mean time, 53

Mercury, Diameter of, 80

" Distance from sun, 80

" Period of, 80

" Rotation of, 80

" Transits of, 81

Metonic cycle, 78

Meyer, Dr., 106

Mimas, satellite of Saturn, 105

Minor planets, 96, 97

Mira Ceti (variable star), 125

Mizar ( $\zeta$  Ursæ Majoris), 117

Moon, Eclipses of, 74

" Diameter of, 61

" Harvest, 68

" Phases of, 65

" Revolution of, 61, 62

" Rotation of, 66

NEAREST fixed star, 123

Nebulae, 130

Neptune, Density of, 109

" Diameter of, 110

" Discovery of, 110

" Distance of, 110

" Satellite of, 110, 111

Newcomb, Prof., 27

Nucleus of comet, 114

Number of visible stars, 26, 27

ONERON, satellite of Uranus, 105

" Offing," 16

Ophiuchi, 70 (binary star), 123

"Opposition" of stars and planets,

33, 92

Orbits of asteroids, 97

Orion, Great nebulae in, 130

"  $\beta$  (Rigel), 29

"  $\epsilon$ , naked eye double, 117

PARABOLA, 113

Pegasi, 85 (binary star), 118

Penumbra in eclipse, 76

Periastron, 122

Perihelion, 46

Persei,  $\beta$  (Algol) (variable star), 126

Phobos, satellite of Mars, 95

Piazzi, 96

Planets, Inferior, 79

" Retrograde motion of, 85, 92

"Plough," 30

Pollux, 30

Procyon, 29

RATIONAL horizon, 22, 23

Regulus ( $\alpha$  Leonis), 32

Rhea, satellite of Saturn, 105

Ricco, Prof., 16

Rigel, 29

Ring, Crape, of Saturn, 105

Rings of Saturn, 105

SAGITTARI,  $\zeta$  (binary star), 123

Saros, cycle, 77

Satellites of planets, 95, 99, 105,

109, 110

Saturn, Density of, 104

" Diameter of, 104

" Distance from sun, 104

" "Dusky ring" of, 105

" Light and heat on, 104

" Period of, 104

" Rotation of, 105

" Satellites of, 105



Schiaparelli, Prof., 80, 83  
 Schur, Dr., 123  
 Seasons, 48  
 Sensible horizon, 22, 23  
 "Sickle" in Leo, 32  
 Sirius, 28  
 " Orbit of, 121  
 Sky visible, A hemisphere, 23, 24  
 Spica ( $\alpha$  Virginis), 29  
 Stars, Binary, 118  
 " Double, 116, 117  
 " Magnitudes of, 28  
 " Nearest, 123  
 " Pole, 29, 30  
 " Variable, 125  
 Sun, Apparent motion of, 32, 33  
 " Diameter of, 57  
 " Distance of, 44, 60  
 " Heat of, 58  
 " Mass of, 57  
 " Reflected image of, 17  
 " Rotation of, 58  
 " Shrinkage of, 58  
 " Specific gravity of, 57  
 " Spots, 58  
 Sun-dial, Time by, 54  
 TAIL of comet, 115  
 Tauri,  $\alpha$  (Aldebaran), 29  
 Temperature, summer and winter,  
 47  
 Tethys, satellite of Saturn, 105

Thomson, Sir William, 58  
 Time, Equation of, 54  
 Titan, satellite of Saturn, 105  
 Titania, satellite of Uranus, 109  
 Tower, Experiment on, 21, 22  
 Transits of Mercury, 81  
 " " Venus, 86, 88  
 Tuttle, Comet of, 114  
 UMBRIEL, satellite of Uranus, 109  
 Uranus, Diameter of, 108  
 " Distance from sun, 108  
 " Period of, 108  
 " Rotation of, 109  
 " Satellites of, 109  
 VARIABLE Stars, 125  
 Vega, ( $\alpha$  Lyrae), 29  
 Venus, Brightness of, 83  
 " Diameter of, 83  
 " Period of, 83  
 " Retrograde motion of, 85  
 " Rotation of, 83  
 " Stationary points, 85  
 " Transits, 86, 88  
 Vesta, minor planet, 96  
 Virginis,  $\alpha$  (Spica), 29  
 "  $\gamma$  (binary star), 122  
 Visible stars, Number of, 26, 27  
 WRUBLEWSKY, 123  
 YOUNG, Prof., 27







ASTRONOMICAL LESSONS - GORE.